

# Vaginal progesterone in women with an asymptomatic sonographic short cervix in the midtrimester decreases preterm delivery and neonatal morbidity: a systematic review and metaanalysis of individual patient data

Roberto Romero, MD; Kypros Nicolaides, MD; Agustin Conde-Agudelo, MD, MPH; Ann Tabor, MD; John M. O'Brien, MD; Elcin Cetingoz, MD; Eduardo Da Fonseca, MD; George W. Creasy, MD; Katharina Klein, MD; Line Rode, MD; Priya Soma-Pillay, MD; Shalini Fusey, MD; Cetin Cam, MD; Zarko Alfirevic, MD; Sonia S. Hassan, MD

**OBJECTIVE:** To determine whether the use of vaginal progesterone in asymptomatic women with a sonographic short cervix ( $\leq 25$  mm) in the midtrimester reduces the risk of preterm birth and improves neonatal morbidity and mortality.

**STUDY DESIGN:** Individual patient data metaanalysis of randomized controlled trials.

**RESULTS:** Five trials of high quality were included with a total of 775 women and 827 infants. Treatment with vaginal progesterone was associated with a significant reduction in the rate of preterm birth  $<33$  weeks (relative risk [RR], 0.58; 95% confidence interval [CI], 0.42–0.80),  $<35$  weeks (RR, 0.69; 95% CI, 0.55–0.88), and  $<28$  weeks (RR, 0.50; 95% CI, 0.30–0.81); respiratory distress syndrome (RR, 0.48; 95% CI, 0.30–0.76); composite neonatal morbidity and mortality

(RR, 0.57; 95% CI, 0.40–0.81); birthweight  $<1500$  g (RR, 0.55; 95% CI, 0.38–0.80); admission to neonatal intensive care unit (RR, 0.75; 95% CI, 0.59–0.94); and requirement for mechanical ventilation (RR, 0.66; 95% CI, 0.44–0.98). There were no significant differences between the vaginal progesterone and placebo groups in the rate of adverse maternal events or congenital anomalies.

**CONCLUSION:** Vaginal progesterone administration to asymptomatic women with a sonographic short cervix reduces the risk of preterm birth and neonatal morbidity and mortality.

**Key words:** admission to neonatal intensive care unit, birthweight  $<1500$  g, mechanical ventilation, prematurity, preterm birth, progesterin, respiratory distress syndrome, transvaginal ultrasound, uterine cervix,  $17\alpha$ -hydroxyprogesterone caproate

Cite this article as: Romero R, Nicolaides K, Conde-Agudelo A, et al. Vaginal progesterone in women with an asymptomatic sonographic short cervix in the midtrimester decreases preterm delivery and neonatal morbidity: a systematic review and metaanalysis of individual patient data. *Am J Obstet Gynecol* 2012;206:124.e1-19.

From the Perinatology Research Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Department of Health and Human Services, Bethesda, MD, and Detroit, MI (Drs Romero, Conde-Agudelo, and Hassan); Department of Obstetrics and Gynecology, King's College Hospital, London, United Kingdom (Dr Nicolaides); Department of Fetal Medicine, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark (Drs Tabor and Rode); Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark (Dr Tabor); Perinatal Diagnostic Center, Central Baptist Hospital and Department of Obstetrics and Gynecology, University of Kentucky, Lexington, KY (Dr O'Brien); Department of Obstetrics and Gynecology, Zeynep Kamil Women and Children Diseases Education and Research Hospital, Uskudar, Istanbul, Turkey (Drs Cetingoz and Cam); Departamento de Obstetrícia e Ginecologia, Hospital do Servidor Público Estadual "Francisco Morato de Oliveira" and School of Medicine, University of São Paulo, São Paulo, Brazil (Dr Fonseca); Columbia Laboratories Inc, Livingston, NJ (Dr Creasy); Department of Obstetrics and Gynecology, Medical University of Vienna, Vienna, Austria (Dr Klein); Department of Obstetrics and Gynecology, Steve Biko Academic Hospital, and the University of Pretoria, Pretoria, South Africa (Dr Soma-Pillay); Department of Obstetrics and Gynecology, Government Medical College and Hospital, Maharashtra, India (Dr Fusey); Department for Women's and Children's Health, University of Liverpool, Liverpool, United Kingdom (Dr Alfirevic); and Department of Obstetrics and Gynecology, Wayne State University/Hutzel Hospital, Detroit, MI (Dr Hassan).

The majority of the authors report no conflict of interest except as stated in this paragraph. J.M.O'B. was involved in studies of progesterone gel treatment for preterm birth prevention sponsored by Columbia Laboratories Inc, the manufacturer of the preparation used in the PREGNANT Trial and a previous trial of vaginal progesterone in women at risk for preterm delivery. J.M.O'B. serves on advisory boards and is a consultant for Watson Pharmaceuticals, a company with a financial interest in marketing vaginal progesterone gel for the prevention of preterm birth. He and others are listed in the patent on the use of all progesterone compounds to prevent preterm birth (US Patent No. 7,884,093: Progesterone for the Treatment and Prevention of Spontaneous Preterm Birth). G.W.C. is an employee of Columbia Laboratories Inc.

This research was supported, in part, by the Intramural Research Program of the Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Department of Health and Human Services.

Reprints not available from the authors.

0002-9378/free • © 2012 Published by Mosby, Inc. • doi: 10.1016/j.ajog.2011.12.003



For Editors' Commentary, see Table of Contents



See related editorial, page 101

Preterm birth is the leading cause of perinatal morbidity and mortality worldwide<sup>1</sup> and contributes to 70% of neonatal mortality and approximately half of long-term neurodevelopmental disabilities.<sup>2</sup> A recent systematic review has estimated that 12.9 million births, or 9.6% of all births worldwide, were preterm, of which approximately 11.9 million (92.3%) were in Africa, Asia, Latin America, and the Caribbean.<sup>3</sup> During the last 25 years, the preterm birth rate in the United States increased 36%, from 9.4% in 1981 to 12.8% in 2006.<sup>4</sup> This increase has been attributed to a higher frequency of “indicated” preterm births in singleton gestations and preterm delivery in multiple gestations resulting, in part, from the use of assisted reproductive technologies.<sup>5-15</sup>

Spontaneous preterm labor/delivery is considered to be one of the “great obstetrical syndromes”,<sup>16,17</sup> a term that emphasizes that obstetrical disorders with a similar phenotype are caused by multiple pathologic processes,<sup>18</sup> have a long subclinical phase, and may result from complex gene-environment interactions.<sup>19-22</sup>

Progesterone is considered a key hormone for pregnancy maintenance, and a decline of progesterone action is implicated in the onset of parturition.<sup>23-26</sup> If such a decline occurs in the midtrimester, cervical shortening may occur, and this would predispose to preterm delivery. Therefore, an untimely decline in progesterone action has been proposed as a mechanism of disease in the “preterm parturition syndrome”.<sup>27</sup>

Progesterone actions are mediated by genomic and nongenomic effects which have been studied in the uterine cervix, myometrium, sperm, etc.<sup>28-101</sup> A blockade of progesterone action can lead to the clinical, biochemical, and morphologic changes associated with cervical ripening.<sup>28-101</sup> A short cervix detected with transvaginal ultrasound is a powerful predictor of preterm birth in women with singleton and twin gestations.<sup>27,102-109</sup> The shorter the sonographic cervical length, the higher the risk of spontaneous preterm birth.<sup>102-105,110-123</sup> Moreover, a short cervix is associated with intraamniotic infection and inflammation, and this may modify the response to interventions.

## ★ EDITORS' CHOICE ★

An interest in the role of progestogens (natural and synthetic) for the prevention of preterm birth has existed for decades.<sup>124-130</sup> Recently, the administration of vaginal progesterone was proposed for the prevention of preterm birth in women with a sonographic short cervix in the midtrimester based on its biologic effects on the cervix, myometrium, and chorioamniotic membranes. In 2007, Fonseca et al,<sup>131</sup> on behalf of the Fetal Medicine Foundation of the United Kingdom, reported that the administration of vaginal progesterone in women with a cervical length  $\leq 15$  mm was associated with a significant 44% reduction in the rate of spontaneous preterm birth  $<34$  weeks of gestation. Similar findings were reported by DeFranco et al<sup>132</sup> in a secondary analysis of a randomized clinical trial of vaginal progesterone in women with a history of preterm birth in which the cervix was measured. Hassan et al<sup>133</sup> reported the largest randomized clinical trial to date, indicating that vaginal progesterone, when administered to women with a cervical length of 10-20 mm, reduces the rate of preterm birth at  $<33$ ,  $<28$ , and  $<35$  weeks, and this was associated with a significant 61% reduction in the rate of respiratory distress syndrome (RDS).<sup>133</sup> Since the publication of the trial of Hassan et al,<sup>133</sup> several trials evaluating vaginal progesterone in women at high risk of spontaneous preterm birth,<sup>134-136</sup> including a subset of women with a short cervix, have been published.

An individual patient data (IPD) metaanalysis is a specific type of systematic review in which the original research data for each participant in a study are sought directly from the investigators responsible for that trial.<sup>137</sup> Such an approach has been considered the gold standard for summarizing evidence across clinical studies since it offers several advantages, both statistically and clinically, over conventional metaanalyses, which are based on published aggregate data.<sup>138</sup> These advantages include standardizing and updating of data sets, the ability to verify the quality of the data and the appropriateness of the analyses, the improvement of consistency across

trials (eg, definition of outcomes), the performance of subgroup analyses that could effectively identify groups of patients who might benefit from an intervention, the investigation of interaction between patient-level covariates and treatment effects, and the performance of time-to-event analyses.<sup>139-141</sup>

Using IPD from randomized controlled trials, we performed a metaanalysis to evaluate the efficacy and safety of vaginal progesterone for the prevention of preterm birth and neonatal morbidity and mortality in asymptomatic women with a sonographic short cervix in the midtrimester. We also sought to determine whether there were clinical benefits associated with the administration of vaginal progesterone in singleton and twin pregnancies.

## MATERIALS AND METHODS

The study was conducted based on a prospectively prepared protocol, and is reported using the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines for metaanalyses of randomized controlled trials<sup>142</sup> and suggested guidelines for IPD metaanalyses.<sup>141</sup>

### Literature search

We searched MEDLINE, EMBASE, CINAHL, and LILACS (all from inception through December 31, 2011); the Cochrane Central Register of Controlled Trials ([www.mrw.interscience.wiley.com/cochrane/cochrane\\_clcentral\\_articles\\_fs.html](http://www.mrw.interscience.wiley.com/cochrane/cochrane_clcentral_articles_fs.html)) (1960 through December 31, 2011); ISI Web of Science ([www.isiknowledge.com](http://www.isiknowledge.com)) (1960 through December 31, 2011); Research Registers of ongoing trials ([www.clinicaltrials.gov](http://www.clinicaltrials.gov), [www.controlled-trials.com](http://www.controlled-trials.com), [www.centerwatch.com](http://www.centerwatch.com), [www.anzctr.org.au](http://www.anzctr.org.au), [www.nih.ac.uk](http://www.nih.ac.uk), and [www.umin.ac.jp/ctr](http://www.umin.ac.jp/ctr)); and Google Scholar using a combination of key words and text words related to *progesterone* (“progesterone,” “progestins,” “progesterone,” “progestagen,” “progestational agent”) and *preterm birth* (“preterm,” “premature”). Proceedings of the Society for Maternal-Fetal Medicine and international meetings on preterm birth, reference lists of identified studies, textbooks, previously published systematic reviews, and review articles were also searched. Experts in the field were contacted to identify further studies. No language restriction was used.

## Study selection

We included randomized controlled trials in which asymptomatic women with a sonographic short cervix (cervical length of  $\leq 25$  mm) in the midtrimester were randomly allocated to receive vaginal progesterone or placebo/no treatment for the prevention of preterm birth. Trials were included if the primary aim of the study was to prevent preterm birth in women with a sonographic short cervix, or if the primary aim was to prevent preterm birth in women with risk factors other than a short cervix, but outcomes were available for patients with a prerandomization cervical length of  $\leq 25$  mm. Trials were excluded if they: (1) were quasirandomized; (2) evaluated vaginal progesterone in women with threatened preterm labor, second trimester bleeding, or premature rupture of membranes; (3) evaluated the administration of vaginal progesterone in the first trimester only to prevent miscarriage; or (4) did not report clinical outcomes. Although there is no agreement on what is a sonographic short cervix, we chose 25 mm as the cutoff because this value corresponds approximately to the 10th percentile for cervical length in the midtrimester.<sup>103,110</sup> In addition, this cervical length is the most commonly used in studies evaluating the predictive accuracy of cervical length for preterm birth.<sup>107,143</sup>

Two investigators (R.R. and A.C.-A.) independently reviewed all potentially relevant articles for eligibility. Disagreements regarding trial eligibility were resolved by consensus.

## Data collection

We contacted the corresponding authors to request access to the data. Authors were asked to supply anonymized data (without identifiers) about patient baseline characteristics, experimental intervention, control intervention, cointerventions, and prespecified outcome measures for every randomly assigned subject and were invited to become part of the collaborative group with joint authorship of the final publication. Data provided by the investigators were merged into a master database specifically constructed for the review. Data were checked for missing information, errors, and inconsistencies by cross-

referencing the publications of the original trials. Quality and integrity of the randomization processes were assessed by reviewing the chronological randomization sequence and pattern of assignment, as well as the balance of baseline characteristics across treatment groups. Inconsistencies or missing data were discussed with the authors and corrections were made when deemed necessary.

## Outcome measures

The prespecified primary outcome measure was preterm birth  $< 33$  weeks of gestation. Secondary outcome measures included preterm birth  $< 37$ ,  $< 36$ ,  $< 35$ ,  $< 34$ ,  $< 30$ , and  $< 28$  weeks of gestation; spontaneous preterm birth  $< 33$  and  $< 34$  weeks of gestation; RDS; necrotizing enterocolitis; intraventricular hemorrhage (all grades); proven neonatal sepsis; retinopathy of prematurity; bronchopulmonary dysplasia; periventricular leukomalacia; fetal death; neonatal death; perinatal mortality, a composite neonatal morbidity and mortality outcome (defined as the occurrence of any of the following events: RDS, intraventricular hemorrhage, necrotizing enterocolitis, proven neonatal sepsis, or neonatal death); Apgar score  $< 7$  at 5 minutes; birthweight  $< 1500$  g and  $< 2500$  g; admission to the neonatal intensive care unit (NICU); use of mechanical ventilation; congenital anomaly; any maternal adverse event; vaginal discharge; vaginal pruritus; discontinuation of treatment because of adverse events; threatened preterm labor; and neurodevelopmental disability at 18-24 months of age. Neonatal morbidities were defined as in the original study.<sup>131,133,134,136,144</sup>

## Assessment of risk of bias


We assessed the risk of bias using the criteria recently outlined in the Cochrane Handbook for Systematic Reviews of Interventions.<sup>137</sup> Seven domains related to risk of bias were assessed in each included trial since there is evidence that these issues are associated with biased estimates of treatment effect: (1) random sequence generation; (2) allocation concealment; (3) blinding of participants and personnel; (4) blinding of outcome

assessment; (5) incomplete outcome data; (6) selective reporting; and (7) other bias. Review authors' judgments were categorized as "low", "high", or "unclear" risk of bias. The assessments considered the risk of *material* bias rather than *any* bias. "Material bias" is defined as a bias of sufficient magnitude to have a notable impact on the results or conclusions of the trial.<sup>137</sup> The risk of bias in each trial included was assessed individually by 2 reviewers (R.R. and A.C.-A.). In addition, methods of random sequence generation, allocation concealment, and blinding were confirmed with the authors of the trials. Any differences of opinion regarding assessment of risk of bias were resolved by discussion.

## Statistical analysis

Statistical analyses were based on an intent-to-treat basis and included all randomized women and their fetuses/infants. For baseline data, maternal outcomes, and gestational age at birth-related outcomes, the unit of analysis was the pregnancy, whereas for neonatal outcomes, the unit of analysis was the neonate. To assess safety of vaginal progesterone, all patients exposed to progesterone were included. This included all studies and patients, even those in which the cervical length was not measured. IPD were combined in a 2-stage approach in which outcomes were analyzed in the original trial and then summary statistics were generated using standard summary data metaanalysis techniques to give an overall measure of effect (summary relative risk [RR] with 95% confidence interval [CI]).<sup>145</sup> Heterogeneity of the results among studies was tested with the quantity  $I^2$ , which describes the percentage of total variation across studies that can be attributed to heterogeneity rather than chance.<sup>146</sup> A value of 0% indicates no observed heterogeneity, whereas  $I^2$  values of  $\geq 50\%$  indicate a substantial level of heterogeneity.<sup>146</sup> We planned to use a fixed effects model if substantial statistical heterogeneity was not present. Random effects models were also used to test the robustness of results. The number needed to treat (NNT) for benefit or harm with the 95% CI was calculated for outcomes for which there was a statistically significant reduc-

**FIGURE 1**  
**Flow of study identification**



Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

tion or increase in risk difference based on control event rates in the trials.<sup>147</sup> Publication and related biases were assessed visually by examining the symmetry of funnel plots and statistically by using the Egger test.<sup>148</sup> A  $P$  value  $< .1$  was considered to indicate significant asymmetry.

Access to data from individual patients also allowed the performance of subgroup analyses to examine whether the administration of vaginal progesterone was more effective in some subgroups than in others. Specifically, we assessed the effect of vaginal progesterone in singleton and twin gestations separately. Also, to explore treatment effects according to other patient characteristics, subgroup analyses were prespecified on the basis of sonographic cervical length ( $<10$ ,  $10$ - $20$ , and  $21$ - $25$  mm), obstetrical history (no previous spontane-

ous preterm birth and at least 1 previous spontaneous preterm birth  $<37$  weeks), maternal age ( $<20$ ,  $20$ - $34$ , and  $\geq 35$  years), race/ethnicity (Caucasian, Black, Asian, and other), and body mass index ( $<18.5$ ,  $18.5$ - $24.9$ ,  $25.0$ - $29.9$ ,  $\geq 30$  kg/m<sup>2</sup>). To explore effects by trial characteristics, prespecified subgroup analyses were planned according to the daily dose of vaginal progesterone ( $90$ - $100$  vs  $200$  mg). Definitions and subgroup analyses were specified before any data were obtained or analyzed. Treatment effects in these subgroups were assessed by simple logistic regression models (which included a subgroup-allocated treatment interaction term), with adjustment for between-trial outcome differences. A test for interaction between treatment and subgroup is the standard method to examine whether treatment effects differ

between subgroups.<sup>149,150</sup> This approach tests and estimates the difference between treatment effects across subgroups directly. It involves one statistical test regardless of the number of subgroups. An interaction  $P$  value  $> .05$  was considered to indicate that the effect of treatment did not differ significantly between subgroups. Adjustment for predictive baseline characteristics, even when largely balanced, can lead to different estimates of treatment effects.<sup>151</sup> Therefore, multivariable logistic regression models were employed to estimate adjusted treatment effects. In the twin pregnancy subgroup, the lack of independence of twins may have influenced the outcome of the analysis. Thus, for adverse perinatal outcomes in twins, we used analytical methods assuming independence between neonates as well as methods recommended to take into account nonindependence of newborns from twin gestations.<sup>137,152</sup> We planned sensitivity analyses to test the robustness of the results by excluding trials with any risk of bias and including only studies for which the primary aim was to assess the effects of vaginal progesterone in women with a short cervix. Subgroup and sensitivity analyses were only performed for the primary outcome of preterm birth  $<33$  weeks of gestation and for the secondary outcome of composite neonatal morbidity and mortality. Analyses were performed with the Review Manager (RevMan) version 5.1 (Nordic Cochrane Centre, Copenhagen, Denmark), and SAS version 9.2 (SAS Institute, Cary, NC) software.

Informed consent was provided by the patients upon enrollment in each of the original trials. In this study, the data were not used for any purposes other than those of the original trial, and no new data were collected. Therefore, informed consent specifically for this project was not considered necessary. No patient identifiers were provided by any investigator. This study was exempted for review by the Human Investigations Committee of Wayne State University's Institutional Review Board, Detroit, MI.

**TABLE 1**  
**Characteristics of studies included**

Study	Participating countries	Primary target population	Inclusion/exclusion criteria	No of women with CL $\leq 25$ mm/fetuses or infants		Intervention	Cointerventions	Primary outcome
				Vaginal progesterone group	Placebo group			
Fonseca et al, <sup>131</sup> 2007	United Kingdom, Chile, Brazil, Greece	Women with a short cervix	Inclusion: women with a singleton or twin pregnancy and sonographic CL $\leq 15$ mm Exclusion: major fetal abnormalities, painful regular uterine contractions, history of ruptured membranes, and cervical cerclage	125/136	125/138	Vaginal progesterone capsule (200 mg/d) or placebo from 24-33 6/7 wk of gestation	Cervical cerclage (1 [0.8%] in vaginal progesterone group and 0 [0.0%] in placebo group)	Spontaneous preterm birth <34 wk
O'Brien et al, <sup>144</sup> 2007	United States, South Africa, India, Czech Republic, Chile, El Salvador	Women with a history of spontaneous preterm birth	Inclusion: women with a singleton pregnancy, gestational age between 16 0/7-22 6/7 wk, and a history of spontaneous singleton preterm birth at 20-35 wk of gestation in the immediately preceding pregnancy Exclusion: planned cervical cerclage, history of adverse reaction to progesterone, treatment with progesterone within 4 wk before enrollment, treatment for seizure disorder, psychiatric illness or chronic hypertension at time of enrollment, history of acute or chronic congestive heart failure, renal failure, uncontrolled diabetes mellitus, active liver disorder, HIV infection with CD4 count of <350 cells/mm <sup>3</sup> and requiring multiple antiviral agents, placenta previa, history or suspicion of breast or genital tract malignancy, history or suspicion of thromboembolic disease, müllerian duct anomaly, major fetal anomaly or chromosomal disorder, or multifetal gestation	12/12	19/19	Vaginal progesterone gel (90 mg/d) or placebo from 18-22 to 37 0/7 wk of gestation, rupture of membranes or preterm delivery, whichever occurred first	None	Preterm birth $\leq 32$ wk
Cetingoz et al, <sup>136</sup> 2011	Turkey	Women at high risk of preterm birth	Inclusion: women with at least 1 previous spontaneous preterm birth, uterine malformation, or twin pregnancy Exclusion: in-place or planned cervical cerclage, serious fetal anomalies	9/14	6/8	Vaginal progesterone suppository (100 mg/d) or placebo from 24-34 wk of gestation	None	Preterm birth <37 wk
Hassan et al, <sup>133</sup> 2011	United States, Republic of Belarus, Chile, Czech Republic, India, Israel, Italy, Russia, South Africa, Ukraine	Women with a short cervix	Inclusion: women with a singleton pregnancy, gestational age between 19 0/7-23 6/7 wk, transvaginal sonographic CL between 10-20 mm, and without signs or symptoms of preterm labor Exclusion: planned cerclage, acute cervical dilation, allergic reaction to progesterone, current or recent progestogen treatment within previous 4 wk, chronic medical conditions that would interfere with study participation or evaluation of treatment, major fetal anomaly or known chromosomal abnormality, uterine anatomic malformation, vaginal bleeding, known or suspected clinical chorioamnionitis	235/235	223/223	Vaginal progesterone gel (90 mg/d) or placebo from 20-23 6/7 to 36 6/7 wk of gestation, rupture of membranes or preterm delivery, whichever occurred first	Emergency cervical cerclage (10 [4.3%] in vaginal progesterone group and 6 [2.7%] in placebo group)	Preterm birth <33 wk

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

(continued)

**TABLE 1**  
**Characteristics of studies included** (continued)

Study	Participating countries	Primary target population	Inclusion/exclusion criteria	No of women with CL $\leq$ 25 mm/fetuses or infants	Vaginal progesterone group	Placebo group	Intervention	Cointerventions	Primary outcome
Rode et al, <sup>1,34</sup> 2011	Denmark, Austria	Women with a twin pregnancy	Inclusion: women with a diamniotic twin pregnancy and chorionicity assessed by ultrasound <16 wk of gestation Exclusion: higher order multiple pregnancies, known allergy to progesterone or peanuts as active treatment contained peanut oil, history of hormone-associated thromboembolic disorders, rupture of membranes, pregnancies treated for or with signs of twin-to-twin transfusion syndrome, intentional fetal reduction, known major structural or chromosomal fetal abnormality, known or suspected malignancy in genitals or breasts, known liver disease	7/14	14/28	Vaginal progesterone pessary (200 mg/d) or placebo from 20-23 6/7 to 33 6/7 wk of gestation	Cervical cerclage (2 [28.6%] in vaginal progesterone group and 2 [14.3%] in placebo group)	Preterm birth <34 wk	

CL, cervical length; HIV, human immunodeficiency virus.

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.


**RESULTS**

**Study selection, details, and quality**

The searches yielded 2611 citations, of which 10 were considered for potential inclusion (Figure 1). Five studies were excluded.<sup>125,153-156</sup> Three of these studies evaluated vaginal progesterone in women at high risk for preterm birth (previous preterm birth,<sup>125,154</sup> uterine malformation,<sup>125</sup> cervical insufficiency,<sup>125</sup> and twins<sup>155</sup>) but none of them measured or collected data on cervical length. Two of these studies<sup>125,154</sup> reported that prophylactic administration of vaginal progesterone reduced the risk of preterm birth in women with a previous preterm birth, whereas the study by Norman et al<sup>155</sup> found that vaginal progesterone did not reduce the risk of the composite outcome delivery or fetal death <34 weeks of gestation in women with a twin gestation. The 2 remaining studies evaluated vaginal progesterone as an adjunct to tocolytic therapy after threatened preterm labor.<sup>153,156</sup> Five studies, which provided data for 775 women (723 [93.3%] with singleton pregnancies and 52 [6.7%] with twin pregnancies) and 827 fetuses/infants (723 [87.4%] from singleton pregnancies and 104 [12.6%] from twin pregnancies), met the inclusion criteria.<sup>131,133,134,136,144</sup>

The main characteristics of studies included in this IPD metaanalysis are shown in Table 1. All studies were double-blind, placebo-controlled trials, of which 4 were multicenter, conducted in sites from both developed and developing countries. Two trials were specifically designed to evaluate the administration of vaginal progesterone in women with a sonographic short cervix,<sup>131,133</sup> one evaluated the use of vaginal progesterone in women with a history of spontaneous preterm birth,<sup>144</sup> another assessed vaginal progesterone in women with a twin gestation,<sup>134</sup> and the remaining trial examined the use of progesterone in women with a prior spontaneous preterm birth, uterine malformations, or twin gestations.<sup>136</sup> Two of these studies<sup>134,144</sup> reported data of planned secondary analyses for women with a short cervix in additional reports.<sup>132,135</sup> Data from the trials by O'Brien et al,<sup>144</sup> Cetinogoz et al,<sup>136</sup> and Rode et al<sup>134</sup> relevant to

**FIGURE 2**  
**Methodological quality summary: risk of biases for each included study**



Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

women with a cervical length of  $\leq 25$  mm before randomization were provided by the authors for inclusion in this review. The 2 trials<sup>131,133</sup> specifically designed to evaluate the use of vaginal pro-

gesterone in women with a short cervix screened a total of 56,711 women, of which 1146 (2.0%) had a sonographic short cervix as defined by the authors for the purposes of each study. Of these

women, 715 (62.4%) were randomized; 708 mothers with their 732 infants provided data for the metaanalysis (~90% of total sample size of the IPD meta-analysis). The other 3 studies provided data for 67 women and 95 infants.


Two studies used vaginal progesterone capsules or pessaries 200 mg/d,<sup>131,134</sup> 2 used vaginal progesterone gel 90 mg/d,<sup>133,144</sup> and the other used vaginal progesterone suppositories 100 mg/d.<sup>136</sup> The treatment was initiated at 24 weeks of gestation in 2 trials,<sup>131,136</sup> between 20-23 weeks of gestation in 2 trials,<sup>133,134</sup> and between 18-22 weeks of gestation in 1 trial.<sup>144</sup> Three studies<sup>131,134,136</sup> reported that participating women received study medication from enrollment until 34 weeks of gestation, and 2 studies<sup>133,144</sup> from enrollment until 36 6/7 weeks of gestation. In 3 studies,<sup>131,133,134</sup> cervical cerclage was allowed after randomization. In the study by Cetingoz et al,<sup>136</sup> cervical cerclage was not performed in any women. The primary outcome was preterm birth <33 weeks of gestation for 1 trial,<sup>133</sup>  $\leq 32$  weeks of gestation for 1 trial,<sup>144</sup> <34 weeks for 1 trial,<sup>134</sup> <37 weeks for 1 trial,<sup>136</sup> and spontaneous preterm birth <34 weeks for the remaining study.<sup>131</sup>

All of the 5 studies included in this IPD metaanalysis had high methodological quality and were considered to be at low risk of bias (Figure 2). One study did not report the method of random sequence generation in the article but reported, upon request, having used a table of random numbers.<sup>131</sup> All 5 studies had adequate allocation concealment and used identical placebo to blind patients and clinical staff to treatment allocation. There was blinding of outcome assessment and adequate handling of incomplete outcome data in all studies. One study<sup>136</sup> did not report several secondary neonatal outcomes of interest to the present study, but they were provided to the investigators (R.R. and A.C.-A.) with the database and included into the meta-analyses. Overall, there was no obvious risk of other biases for the 5 trials.

**Primary outcome**

Treatment with vaginal progesterone in patients with a sonographic short cervix

**FIGURE 3**  
Effect of vaginal progesterone on preterm birth <33 weeks of gestation



Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

was associated with a significant reduction in the risk of preterm birth <33 weeks of gestation (12.4% vs 22.0%; RR, 0.58; 95% CI, 0.42–0.80;  $I^2 = 0\%$ ; 775 women) (Figure 3). The number of patients with a short cervix who needed to be treated with vaginal progesterone rather than with placebo to prevent 1 case of preterm birth <33 weeks of gestation was 11 (95% CI, 8–23).

### Secondary outcomes

Patients allocated to receive vaginal progesterone had a significantly lower risk of preterm birth <35 weeks of gestation (20.4% vs 30.5%; RR, 0.69; 95% CI, 0.55–0.88;  $I^2 = 0\%$ ; NNT for benefit 11; 95% CI, 7–27), <34 weeks (16.0% vs 27.1%; RR, 0.61; 95% CI, 0.47–0.81;  $I^2 = 0\%$ ; NNT for benefit 9; 95% CI, 7–19), <30 weeks (7.5% vs 13.2%; RR, 0.58; 95% CI, 0.38–0.89;  $I^2 = 0\%$ ; NNT for benefit 18; 95% CI, 12–69), and <28 weeks (5.4% vs 11.1%; RR, 0.50; 95% CI, 0.30–0.81;  $I^2 = 0\%$ ; NNT for benefit 18; 95% CI, 13–47) compared to those allocated to placebo (Table 2). Moreover, vaginal progesterone administration was associated with a significantly reduced risk of spontaneous preterm birth <33 and <34 weeks of gestation. The reduction in the risk of preterm birth <36 weeks of gestation was marginally significant (RR, 0.82; 95% CI, 0.67–1.00).

Treatment with vaginal progesterone was associated with an overall nonsignificant reduction in the risk of perinatal mortality (3.4% vs 5.3%; RR, 0.63; 95% CI, 0.34–1.18;  $I^2 = 41\%$ ). This reduction appeared to be attributable to a reduction in neonatal death (1.9% vs 3.6%; RR, 0.55; 95% CI, 0.26–1.19;  $I^2 = 43\%$ ) rather than fetal death (1.5% vs 1.7%; RR, 0.82; 95% CI, 0.28–2.42;  $I^2 = 0\%$ ).

Infants whose mothers received vaginal progesterone also had a significantly lower risk of RDS (6.1% vs 12.5%; RR, 0.48; 95% CI, 0.30–0.76;  $I^2 = 0\%$ ; NNT for benefit 15; 95% CI, 11–33), composite neonatal morbidity and mortality (9.7% vs 17.3%; RR, 0.57; 95% CI, 0.40–0.81;  $I^2 = 0\%$ ; NNT for benefit 13; 95% CI, 10–30), birthweight <1500 g (8.8% vs 16.5%; RR, 0.55; 95% CI, 0.38–0.80;  $I^2 = 6\%$ ; NNT for benefit 13; 95% CI, 10–30), admission to NICU (20.7% vs 29.1%; RR, 0.75; 95% CI, 0.59–0.94;  $I^2 = 0\%$ ; NNT for benefit 14; 95% CI, 8–57), and mechanical ventilation (8.5% vs 12.3%; RR, 0.66; 95% CI, 0.44–0.98;  $I^2 = 0\%$ ; NNT for benefit 24; 95% CI, 15–408) than infants whose mothers had received placebo.

There was no evidence of an effect of vaginal progesterone on necrotizing enterocolitis, intraventricular hemorrhage, proven neonatal sepsis, retinopathy of prematurity, bronchopulmonary dysplasia, periventricular leukomalacia, Ap-

gar score <7 at 5 minutes, birthweight <2500 g, or threatened preterm labor.

In addition, the rates of maternal adverse effects, discontinuation of treatment because of adverse effects, and congenital anomalies did not differ significantly between the vaginal progesterone and placebo groups. One study<sup>134</sup> reported that the mean Ages and Stages Questionnaire scores (a tool that measures neurodevelopmental disability) at 18 months of age were  $193 \pm 42.6$  for infants in the progesterone group and  $194 \pm 40.6$  for infants in the placebo group ( $P = .89$ ).

### Effect of vaginal progesterone in singleton and twin gestations

Table 3 shows the effect of vaginal progesterone on the risk of preterm birth and perinatal outcomes in singleton and twin gestations separately. According to the interaction  $P$  values (all > .10), there was no evidence that women with singleton pregnancies benefit more or less from the use of vaginal progesterone than women with twin pregnancies.

Among singleton gestations, the administration of vaginal progesterone was associated with a statistically significant reduction in the risk of preterm birth <33, <35, and <28 weeks of gestation; RDS; composite neonatal morbidity and mortality; Apgar score <7 at 5 minutes; birthweight <1500 g; and admission to the NICU.

Among twin gestations, the administration of progesterone did not significantly reduce the risk of preterm birth <33 weeks of gestation (RR, 0.70; 95% CI, 0.34–1.44). However, it significantly decreased the risk of composite neonatal morbidity and mortality (RR, 0.52; 95% CI, 0.29–0.93). There were no significant differences in other outcome measures among the vaginal progesterone and placebo groups. The effects of vaginal progesterone on adverse perinatal outcome in twins were tested with 2 methods: one that assumed independence of the twins, and another that did not make such assumption. The results of both analyses were similar, with a slightly wider CI when the method which assumed nonindependence was employed (Table 4). The beneficial effects on composite neonatal morbidity and mortality in twins remained statistically significant (RR, 0.56; 95% CI, 0.30–0.97).

**TABLE 2**  
**Effect of vaginal progesterone on secondary outcome measures<sup>a</sup>**

Outcome	No. of trials	No. of events/total no.		Pooled RR (95% CI)	I <sup>2</sup> (%)	NNT (95% CI)
		Vaginal progesterone	Placebo			
Preterm birth <37 wk	5	144/388	165/387	0.89 (0.75–1.06)	0	–
Preterm birth <36 wk	5	108/388	136/387	0.82 (0.67–1.00)	0	–
Preterm birth <35 wk	5	79/388	118/387	0.69 (0.55–0.88)	0	11 (7–27)
Preterm birth <34 wk	5	62/388	105/387	0.61 (0.47–0.81)	0	9 (7–19)
Preterm birth <30 wk	5	29/388	51/387	0.58 (0.38–0.89)	0	18 (12–69)
Preterm birth <28 wk	5	21/388	43/387	0.50 (0.30–0.81)	0	18 (13–47)
Spontaneous preterm birth <33 wk	5	39/388	71/387	0.57 (0.40–0.81)	0	13 (9–29)
Spontaneous preterm birth <34 wk	5	51/388	87/387	0.62 (0.46–0.84)	0	12 (8–28)
Respiratory distress syndrome	5	25/411	52/416	0.48 (0.30–0.76)	0	15 (11–33)
Necrotizing enterocolitis	5	5/411	6/416	0.88 (0.30–2.64)	0	–
Intraventricular hemorrhage	5	6/411	9/416	0.74 (0.27–2.05)	0	–
Proven neonatal sepsis	5	12/411	20/416	0.64 (0.32–1.29)	13	–
Retinopathy of prematurity	5	6/411	3/416	1.56 (0.46–5.28)	0	–
Bronchopulmonary dysplasia	2	4/249	5/231	0.76 (0.21–2.79)	NA	–
Periventricular leukomalacia	2	0/249	0/231	Not estimable	NA	–
Fetal death	5	6/411	7/416	0.82 (0.28–2.42)	0	–
Neonatal death	5	8/411	15/416	0.55 (0.26–1.19)	43	–
Perinatal death	5	14/411	22/416	0.63 (0.34–1.18)	41	–
Composite neonatal morbidity/mortality <sup>a</sup>	5	40/411	72/416	0.57 (0.40–0.81)	0	13 (10–30)
Apgar score <7 at 5 min	5	15/408	27/412	0.57 (0.32–1.02)	16	–
Birthweight <1500 g	5	36/410	68/413	0.55 (0.38–0.80)	6	13 (10–30)
Birthweight <2500 g	5	140/410	162/413	0.91 (0.76–1.08)	0	–
Admission to NICU	5	85/411	121/416	0.75 (0.59–0.94)	0	14 (8–57)
Mechanical ventilation	5	35/411	51/416	0.66 (0.44–0.98)	0	24 (15–408)
Congenital anomaly	7	30/1967	34/1954	0.89 (0.55–1.44)	0	–
Any maternal adverse event	3	86/624	80/595	1.04 (0.79–1.38)	0	–
Vaginal discharge	4	244/1065	248/1057	1.00 (0.87–1.15)	33	–
Vaginal pruritus	4	54/1065	50/1057	1.08 (0.74–1.57)	0	–
Discontinuation of treatment because of adverse events	5	28/1083	28/1061	1.01 (0.61–1.69)	0	–
Threatened preterm labor	5	115/384	139/383	0.83 (0.68–1.02)	16	–
Low ASQ developmental and socioemotional score at 18 mo of age <sup>b</sup>	1	19/503	18/488	1.02 (0.54–1.93)	NA	–

ASQ, Ages and Stages Questionnaire; CI, confidence interval; NA, not applicable; NICU, neonatal intensive care unit; NNT, number needed to treat; RR, relative risk.

<sup>a</sup> Occurrence of any of the following events: respiratory distress syndrome, intraventricular hemorrhage, necrotizing enterocolitis, proven neonatal sepsis, or neonatal death; <sup>b</sup> ASQ score <115 points.

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

Importantly, vaginal progesterone was associated with a significant reduction in the risk of preterm birth <33 weeks of gestation in both women with a singleton gestation

with no previous preterm birth (RR, 0.60; 95% CI, 0.39–0.92) as well as in women with a singleton gestation and at least 1 previous spontaneous preterm birth <37 weeks of

gestation (RR, 0.54; 95% CI, 0.30–0.98). Moreover, vaginal progesterone significantly decreased the risk of composite neonatal morbidity and mortality in women with a

TABLE 3

## Effect of vaginal progesterone on preterm birth and perinatal outcomes in singleton and twin gestations

Outcome	Singleton pregnancy				Twin pregnancy				Interaction P value
	No. of trials	No. of events/total No.		Pooled RR (95% CI)	No. of trials	No. of events/total No.		Pooled RR (95% CI)	
		Vaginal progesterone	Placebo			Vaginal progesterone	Placebo		
Primary outcome									
Preterm birth <33 wk	4	41/365	72/358	0.56 (0.40–0.80)	3	7/23	13/29	0.70 (0.34–1.44)	.55
Secondary outcomes									
Preterm birth <37 wk	4	127/365	141/358	0.91 (0.75–1.10)	3	17/23	24/29	0.91 (0.68–1.23)	.88
Preterm birth <35 wk	4	67/365	100/358	0.67 (0.51–0.87)	3	12/23	18/29	0.91 (0.57–1.46)	.24
Preterm birth <28 wk	4	20/365	39/358	0.51 (0.31–0.85)	3	1/23	4/29	0.44 (0.11–1.85)	.83
Respiratory distress syndrome	4	17/365	37/358	0.47 (0.27–0.81)	3	8/46	15/58	0.48 (0.21–1.09)	.68
Necrotizing enterocolitis	4	5/365	6/358	0.88 (0.29–2.62)	3	0/46	0/58	Not estimable	NA
Intraventricular hemorrhage	4	5/365	7/358	0.68 (0.22–2.13)	3	1/46	2/58	1.00 (0.10–10.11)	.74
Proven neonatal sepsis	4	11/365	14/358	0.80 (0.37–1.74)	3	1/46	6/58	0.33 (0.06–1.67)	.30
Retinopathy of prematurity	4	5/365	3/358	1.51 (0.40–5.69)	3	1/46	0/58	1.42 (0.05–42.22)	.91
Fetal death	4	6/365	7/358	0.82 (0.28–2.40)	3	0/46	0/58	Not estimable	NA
Neonatal death	4	6/365	11/358	0.53 (0.20–1.39)	3	2/46	4/58	0.68 (0.23–2.02)	.69
Perinatal death	4	12/365	18/358	0.64 (0.31–1.31)	3	2/46	4/58	0.68 (0.23–2.02)	.90
Composite neonatal morbidity/mortality <sup>a</sup>	4	29/365	49/358	0.59 (0.38–0.91)	3	11/46	23/58	0.52(0.29–0.93)	.69
Apgar score <7 at 5 min	4	11/362	23/354	0.48 (0.24–0.95)	3	4/46	4/58	1.03 (0.38–2.81)	.20
Birthweight <1500 g	4	28/364	53/355	0.52 (0.34–0.81)	3	8/46	15/58	0.69 (0.34–1.39)	.47
Birthweight <2500 g	4	102/364	117/355	0.86 (0.69–1.07)	3	38/46	45/58	1.11 (0.92–1.35)	.11
Admission to NICU	4	59/365	87/358	0.67 (0.50–0.91)	3	26/46	34/58	0.98 (0.70–1.35)	.12
Mechanical ventilation	4	28/365	43/358	0.65 (0.41–1.01)	3	7/46	8/58	0.68 (0.30–1.56)	.88

CI, confidence interval; NA, not applicable; NICU, neonatal intensive care unit; RR, relative risk.

<sup>a</sup> Occurrence of any of the following events: respiratory distress syndrome, intraventricular hemorrhage, necrotizing enterocolitis, proven neonatal sepsis, or neonatal death.

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

singleton gestation and at least 1 previous spontaneous preterm birth <37 weeks of gestation (RR, 0.41; 95% CI, 0.17–0.98), and in women with a twin gestation and no previous preterm birth (RR, 0.52; 95% CI, 0.29–0.93).

### Subgroup and sensitivity analyses

Subgroup analyses of the effect of vaginal progesterone on primary outcomes are presented in Table 5. There was no evidence that women in any one of the prespecified subgroups benefit more or less from the use of vaginal progesterone than those in any other subgroup (all *P* for interaction > .30). However, the use of vaginal progesterone was associated with a statistically significant reduction in the risk of preterm birth <33 weeks and composite neonatal morbidity and

mortality in both women with no previous spontaneous preterm birth and women with at least 1 previous spontaneous preterm birth <37 weeks of gestation, women with a sonographic cervical length between 10–20 mm, women aged 20–34 years, and Caucasian women.

No significant differences were noted for preterm birth <33 weeks of gestation and composite neonatal morbidity and mortality between subgroups based on the daily dose of progesterone. A significant decrease in the risk of preterm birth <33 weeks of gestation and composite neonatal morbidity and mortality was found in women who received either 90–100 or 200 mg/d of vaginal progesterone.

The effect of vaginal progesterone on the risk of preterm birth <33 weeks of

gestation and composite neonatal morbidity/mortality did not change when sensitivity analysis was limited to the 2 trials<sup>131,133</sup> in which the primary aim was to evaluate the effect of vaginal progesterone in women with a short cervix (pooled RR, 0.57; 95% CI, 0.40–0.80 for preterm birth <33 weeks and pooled RR, 0.54; 95% CI, 0.35–0.82 for composite neonatal morbidity/mortality). In addition, the results of the metaanalyses did not change significantly when random effects models were used for preterm birth <33 weeks (RR, 0.59; 95% CI, 0.43–0.81) or for composite neonatal morbidity/mortality (RR, 0.59; 95% CI, 0.41–0.83). Sensitivity analyses based on trial quality were not performed because all trials were considered at low risk for

biases. No funnel plots showed asymmetry, either visually or in terms of statistical significance ( $P > .10$  for all, by Egger test).

**COMMENT**

**Principal findings of this study**

Vaginal progesterone administration to asymptomatic women with a sonographic short cervix in the midtrimester was associated with: (1) a significant 42% reduction in the rate of preterm birth <33 weeks (primary outcome); (2) a significant reduction in the risk of preterm birth <35, <34, <30, and <28 weeks and a trend for a reduction in the rate of preterm birth <36 weeks; (3) a significant reduction in the risk of spontaneous preterm birth <33 and <34 weeks; (4) a significantly lower rate of RDS (6.1% vs 12.5% in the placebo group); (5) a significant 43% decrease in composite neonatal morbidity and mortality; (6) a significantly lower rate of admission to NICU (20.7% vs 29.1%) and use of mechanical ventilation (8.5% vs 12.3%); (7) a significantly lower rate of neonates with a birthweight <1500 g (8.8% vs 16.5%); and (8) a nonsignificant difference in the rate of maternal adverse events (13.8% vs 13.4%), discontinuation of therapy because of adverse events (2.6% vs 2.6%), congenital anomalies (1.5% vs 1.7%), and neurodevelopmental disability at 18 months of age (3.8% vs 3.7%). Furthermore, most results remained significant when the analyses were restricted to patients with a singleton gestation. In patients with a twin gestation, there was a nonsignificant trend toward reduction of the rate of preterm birth <33 weeks of gestation. However, there was a significant reduction in the risk of composite neonatal morbidity/mortality (pooled RR, 0.52; 95% CI, 0.29–0.93). Importantly, the reduction in the rates of preterm birth <33 weeks of gestation and composite neonatal morbidity and mortality was observed in both women with no previous spontaneous preterm birth and women with a history of spontaneous preterm birth. Finally, there was no difference in efficacy when a dose of either 90-100 or 200 mg/d of vaginal progesterone was used.

The major effect of vaginal progesterone was to reduce the rate of early pre-

**TABLE 4**  
**Effect of vaginal progesterone on adverse perinatal outcomes in twins according to analytical method used**

Outcome	Pooled RR (95% CI)	
	Assuming independence of newborns	Adjustment for the lack of independence of newborns
Respiratory distress syndrome	0.48 (0.21–1.09)	0.58 (0.25–1.39)
Necrotizing enterocolitis	Not estimable	Not estimable
Intraventricular hemorrhage	1.00 (0.10–10.11)	1.00 (0.05–18.19)
Proven neonatal sepsis	0.33 (0.06–1.67)	0.44 (0.04–4.67)
Retinopathy of prematurity	1.42 (0.05–42.22)	1.36 (0.03–58.74)
Fetal death	Not estimable	Not estimable
Neonatal death	0.68 (0.23–2.02)	0.48 (0.06–3.74)
Perinatal death	0.68 (0.23–2.02)	0.48 (0.06–3.74)
Composite neonatal morbidity/mortality <sup>a</sup>	0.52 (0.29–0.93)	0.56 (0.30–0.97)
Apgar score <7 at 5 min	1.03 (0.38–2.81)	0.88 (0.16–4.75)
Birthweight <1500 g	0.69 (0.34–1.39)	0.73 (0.29–1.83)
Birthweight <2500 g	1.11 (0.92–1.35)	1.13 (0.91–1.40)
Admission to NICU	0.98 (0.70–1.35)	0.89 (0.60–1.31)
Mechanical ventilation	0.68 (0.30–1.56)	0.60 (0.22–1.65)

CI, confidence interval; NICU, neonatal intensive care unit; RR, relative risk.

<sup>a</sup> Occurrence of any of the following events: respiratory distress syndrome, intraventricular hemorrhage, necrotizing enterocolitis, proven neonatal sepsis, or neonatal death.

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

term birth; however, our results indicate that a fraction of late preterm births (34-36 6/7 weeks) can also be prevented with the administration of vaginal progesterone. Further studies are required to explore the reasons for the differential effect in early vs late preterm births. One possibility is that vaginal progesterone was stopped at 36 6/7 weeks of gestation in the largest trial, but at 34 weeks of gestation in the trial of Fonseca et al.<sup>131</sup> Nonetheless, the importance of preventing early preterm birth stems from their disproportionate contribution to serious perinatal morbidity and long-term neurodevelopmental disability.

A decrease in the rate of preterm birth has been considered a surrogate endpoint for neonatal morbidity and mortality, and indeed, in some trials, a reduction in the rate of preterm birth has not been accompanied by a demonstrable reduction in the frequency of neonatal morbid events. It

has been argued that a preventive strategy for preterm birth should accomplish both a reduction in preterm birth and neonatal morbidity. Hassan et al<sup>133</sup> demonstrated that the significant reduction in the rate of preterm birth at <33 weeks was associated with a significant reduction in the rate of RDS by 61%, whereas Fonseca et al<sup>131</sup> reported a nonsignificant reduction in the risk of RDS by 41%. However, in this IPD metaanalysis, we found that vaginal progesterone significantly decreased the risk of RDS by 52%.

There was no significant difference in the risk of adverse maternal events, discontinuation of treatment because of adverse events, and congenital anomalies between vaginal progesterone and placebo groups. Only 1 study of twin gestations<sup>134</sup> examined developmental and socioemotional scores at 18 months of age, and found that there was no difference between infants ex-

TABLE 5

Subgroup analyses of effect of vaginal progesterone on preterm birth <33 weeks of gestation and composite neonatal morbidity/mortality<sup>a</sup>

Subgroup	Preterm birth <33 wk of gestation			Composite neonatal morbidity/mortality <sup>a</sup>		
	n	RR (95% CI)	Interaction P value	n	RR (95% CI)	Interaction P value
Patient characteristics						
Cervical length, mm			.32			.93
<10	79	0.83 (0.49–1.41)		90	0.62 (0.28–1.38)	
10-20	653	0.52 (0.35–0.76)		680	0.54 (0.35–0.84)	
21-25	43	0.50 (0.10–2.41)	.68	57	0.55 (0.26–1.19)	.40
Obstetric history						
With no previous preterm birth	606	0.61 (0.42–0.89)		658	0.62 (0.43–0.91)	
With ≥1 previous preterm birth	169	0.54 (0.30–0.98)		169	0.41 (0.17–0.96)	
Maternal age, y			.85			.31
<20	63	0.66 (0.21–2.14)		66	1.05 (0.25–4.37)	
20-34	620	0.58 (0.41–0.84)		659	0.48 (0.31–0.73)	
≥35	92	0.49 (0.20–1.15)		102	0.89 (0.33–2.36)	
Race/ethnicity			.44			.68
Caucasian	269	0.39 (0.22–0.69)		291	0.57 (0.35–0.93)	
Black	287	0.74 (0.46–1.19)		293	0.60 (0.32–1.12)	
Asian	157	0.53 (0.21–1.34)		159	0.87 (0.20–3.78)	
Other	41	0.60 (0.19–1.92)		42	0.20 (0.03–1.57)	
Body mass index, kg/m <sup>2</sup>			.70			.58
<18.5	58	0.35 (0.10–1.20)		62	0.26 (0.05–1.34)	
18.5-24.9	359	0.63 (0.36–1.10)		390	0.62 (0.37–1.03)	
25.0-29.9	187	0.68 (0.39–1.19)		200	0.76 (0.39–1.47)	
≥30	159	0.49 (0.26–0.92)		163	0.46 (0.21–1.03)	
Trial characteristics						
Daily dose of vaginal progesterone, mg			.57			.92
90-100	504	0.53 (0.33–0.85)		511	0.58 (0.35–0.95)	
200	271	0.63 (0.41–0.96)		316	0.56 (0.34–0.94)	

CI, confidence interval; RR, relative risk.

<sup>a</sup> Occurrence of any of the following events: respiratory distress syndrome, intraventricular hemorrhage, necrotizing enterocolitis, proven neonatal sepsis, or neonatal death.

Romero. Vaginal progesterone to prevent preterm birth in women with a short cervix: an IPD metaanalysis. *Am J Obstet Gynecol* 2012.

posed to vaginal progesterone and those exposed to placebo. These results are consistent with those of an unpublished observation in a trial of singleton gestations exposed to vaginal progesterone.<sup>157</sup>

### Subgroup analyses

Subgroup analyses did not indicate that vaginal progesterone has differential efficacy in the main clinical subgroups of interest. For example, patients with a sonographic short cervix with or without a

history of preterm birth seem to benefit from vaginal progesterone for the reduction of preterm birth. On the other hand, there was some suggestion that patients with a singleton gestation, sonographic cervical length between 10-20 mm, history of preterm birth, age 20-34 years, Caucasian, and body mass index ≥30 kg/m<sup>2</sup> might derive a larger benefit from the use of vaginal progesterone than those with other characteristics. Although such anal-

ysis was prespecified, subgroup analyses should, of course, be interpreted cautiously because of the risk for false-positive and false-negative results.<sup>149,150</sup>

An important question is the range of cervical length in which vaginal progesterone is effective. Fonseca et al<sup>131</sup> noted that vaginal progesterone reduced the rate of spontaneous preterm delivery at <34 weeks of gestation by only 15% in women with a cervical length of 1-5 mm

and 25% in patients with a cervical length of 6–10 mm, but the effect size was 75% in patients with a cervical length between 11–15 mm. One possible explanation for this observation is that women with a very short cervix are more likely to have intraamniotic inflammation and may be less responsive to progesterone.<sup>158,159</sup> These observations are in keeping with the rationale for excluding patients with a cervical length of <10 mm in the study reported by Hassan et al.<sup>133</sup> A subgroup analysis of this IPD metaanalysis suggests that progesterone might be less effective in patients with a cervical length of <10 mm. However, there was no statistically significant differential effect according to cervical length (interaction *P* value of .32 for preterm birth <33 weeks of gestation and .93 for composite neonatal morbidity/mortality). Therefore, this result is subject to all limitations of analysis of subgroups.<sup>149,150</sup>

Fonseca et al<sup>131</sup> and Hassan et al<sup>133</sup> reported that vaginal progesterone was associated with a nonstatistically significant reduction in the rate of preterm birth <34 weeks and preterm birth <33 weeks, respectively, in patients with a short cervix and history of spontaneous preterm birth. Some have interpreted such findings as suggesting that vaginal progesterone does not reduce the rate of preterm birth in these women. This is not a correct interpretation of the results of the trials, because the number of patients with a prior preterm birth in each individual trial was small. Indeed, patients with a history of preterm birth <37 weeks of gestation represented only 15% and 21% of those enrolled in the trials of Fonseca et al<sup>131</sup> and Hassan et al,<sup>133</sup> respectively. Moreover, the primary objective of both trials was to test whether vaginal progesterone would reduce the rate of preterm birth in women with a sonographic short cervix, and not in a particular subgroup. Although the inclusion of patients with a history of spontaneous preterm birth was not a central focus in the design of these trials, this IPD metaanalysis sheds light on this question, because with the increased statistical power afforded by the larger sample size, we were able to show that patients with a short cervix and a history of

preterm birth benefitted to the same extent as patients with a short cervix without a history of preterm birth. There is no biological explanation as to why patients with a history of spontaneous preterm birth would not benefit from progesterone administration if they have a short cervix.

The results reported herein have clinical implications because they extend the indication of vaginal progesterone to women with a history of spontaneous preterm birth with a short cervix. Some have claimed that 17 $\alpha$ -hydroxyprogesterone caproate is the only therapeutic intervention effective in reducing the rate of preterm birth in women with a history of preterm birth<sup>160</sup>—such conclusion is contradicted by the results of this IPD metaanalysis. In addition, a recent metaanalysis reported that cervical cerclage (compared with no cervical cerclage) significantly reduces the risk of preterm birth <35 weeks of gestation by 30% (RR, 0.70; 95% CI, 0.55–0.89) and composite perinatal morbidity and mortality by 36% (RR, 0.64; 95% CI, 0.45–0.91) in women with a singleton gestation, previous spontaneous preterm birth, and cervical length <25 mm.<sup>161</sup> In the present IPD metaanalysis, vaginal progesterone significantly decreased the risk of preterm birth <33 weeks of gestation by 46% (RR, 0.54; 95% CI, 0.30–0.98) in patients with a singleton gestation, prior preterm birth, and sonographic short cervix. Therefore, it appears that administration of vaginal progesterone could be an alternative treatment to cervical cerclage in patients with a singleton pregnancy, short cervix, and history of spontaneous preterm birth for preventing preterm birth and neonatal morbidity and mortality. Vaginal progesterone administration does not carry the risks of anesthesia, the surgical procedure per se, or some of the complications attributed to cerclage (ie, rupture of membranes).<sup>162–171</sup>

The role of progestins in women with a twin gestation has been of interest to several investigators. Several randomized clinical trials have evaluated the effects of 17 $\alpha$ -hydroxyprogesterone caproate or vaginal progesterone for the prevention of preterm birth in twin gestations, and

the results have been uniformly negative.<sup>126,172–174</sup> However, the excess rate of preterm birth of twin gestations is due to multiple causes. In both singleton and multiple gestations, preterm labor is syndromic; therefore, it is unrealistic to expect that one treatment will reduce the rate of preterm birth in all cases.<sup>175</sup> Thus, we explored the hypothesis that vaginal progesterone may benefit women with twin gestations and a short cervix. This IPD metaanalysis revealed a 30% non-significant reduction in the rate of preterm birth <33 weeks of gestation (30.4% vs 44.8%; RR, 0.70; 95% CI, 0.34–1.44). Importantly, vaginal progesterone was associated with a significant reduction in composite neonatal morbidity and mortality (23.9% vs 39.7%; RR, 0.52; 95% CI, 0.29–0.93). We believe that a randomized controlled trial is urgently needed to explore whether women with dichorionic twin gestations and a short cervix may benefit from vaginal progesterone.

Randomized controlled trials included in this IPD metaanalysis have used 3 different doses and formulations of vaginal progesterone: progesterone gel with 90 mg, progesterone suppositories with 100 mg, and progesterone suppositories with 200 mg. To explore whether the dose altered the effectiveness of treatment, we conducted a subgroup analysis comparing patients allocated to receive 90–100 mg/d vs those who received 200 mg/d. Both doses were associated with a statistically significant reduction in the rate of preterm birth <33 weeks and composite neonatal morbidity and mortality. Yet, the only primary trial that showed a reduction in preterm birth, RDS, and composite morbidity was that of Hassan et al,<sup>133</sup> which used 90 mg daily. This represents level-1 evidence of efficacy. The findings of this IPD metaanalysis favor the use of a daily vaginal administration of 90 mg of progesterone because it is the lowest dose that reduced the risk of preterm birth <33 weeks and neonatal morbidity and mortality. Patients who used 90 mg/d of vaginal progesterone received it in a gel, whereas patients who used either 100 or 200 mg/d of vaginal progesterone received it in a suppository. It is known that these suppositories melt in the vagina and there is often loss of the product

over the course of a day. The gel is administered as a bioadhesive preparation applied against the vaginal wall; therefore, it is less likely to lead to loss of the active compound.

### Strengths and limitations

The reliability and robustness of the results reported herein are supported by several considerations. First, the access to data from individual patients enabled a more rigorous analysis than what is possible from published data. The collection of data from individual patients allowed the use of previously unreported data, improved assessment of the study quality, standardization of outcome measures, undertaking an intent-to-treat analysis, and use of optimal analytical methods. Subgroup and multivariable analyses would not have been possible without the availability of individual patient data. Second, a rigorous methodology for performing a systematic review and IPD metaanalysis of randomized controlled trials was employed. Third, we retrieved data for most patients with a sonographic short cervix included in randomized controlled trials of vaginal progesterone. We obtained data for 775 patients with a sonographic cervical length  $\leq 25$  mm from 5 studies. Data for approximately 30 women with a cervical length  $\leq 25$  mm from 3 studies that did not measure or collect data on cervical length were not available for our metaanalysis. Thus, we were able to retrieve individual data from at least 96% of patients with a sonographic short cervix who were randomized to receive vaginal progesterone or placebo. Since a large proportion of data were obtained, the results are likely to be representative. Fourth, the methodological quality of all trials included in the review was high. Fifth, there was evidence of clinical and statistical homogeneity for the primary outcome and for most of the secondary outcomes evaluated. Sixth, subgroup analyses were performed according to patient characteristics at trial entry and trial characteristics. Finally, the sensitivity analyses were consistent with the primary results.

The study has limitations. First, some subgroup analyses were based on a small

number of patients. As a result, the analyses were limited in their power to detect differences, if any existed. Second, 3 trials evaluated the effect of vaginal progesterone in women at high risk for preterm delivery, but the investigators did not measure the cervix or the data were not collected. However, it is unlikely that the overall estimate of effect size in our study would change with the inclusion of approximately 30 patients with a short cervix from such studies. Third, to date, only 1 study<sup>134</sup> has reported on the neurodevelopmental outcomes of children at 18 months of age. Another report in abstract form suggests that there is no evidence of adverse outcome at 24 months of age.<sup>157</sup> Collectively, these observations would be consistent with the long-standing view that the administration of progesterone during pregnancy is safe. Such a view is derived from studies in which vaginal progesterone is used in the first trimester of pregnancy of patients undergoing assisted reproductive technologies.

Another limitation is that neonatal morbidity was not collected consistently across the studies. For example, the study of Hassan et al<sup>133</sup> collected information about bronchopulmonary dysplasia, but the study of Fonseca et al<sup>131</sup> did not. Similarly, some studies did not collect information about the intraventricular hemorrhage grade.<sup>134,136</sup>

### Cost-effectiveness of the intervention

Thus far, 2 studies have evaluated the cost-effectiveness of routine transvaginal cervical length measurement and treatment with vaginal progesterone to prevent preterm birth and resultant neonatal morbidity and mortality. In 2010, Cahill et al<sup>176</sup> reported that a strategy of universal cervical length screening at the time of the routine fetal anatomy sonogram to identify women with a cervical length of  $\leq 15$  mm and subsequent treatment with vaginal progesterone was the most cost-effective strategy and the dominant choice over the following 3 alternatives: cervical length screening for women at increased risk for preterm birth and treatment with vaginal progesterone, risk-based treatment with 17 $\alpha$ -hydroxyprogesterone caproate without

screening, and no screening or treatment. These investigators concluded that universal screening of cervical length and treatment with vaginal progesterone would be the most effective of the different approaches considered. Recently, Werner et al<sup>177</sup> compared the cost-effectiveness of 2 strategies, no routine cervical length screening and single routine transvaginal cervical length measurement at 18-24 weeks of gestation followed by treatment with vaginal progesterone if cervical length  $< 15$  mm. This study showed that routine cervical length screening/use of vaginal progesterone was the dominant strategy when compared to routine care. For every 100,000 women screened, 22 cases of neonatal death or long-term neurologic deficits could be prevented, and approximately \$19 million could potentially be saved. Therefore, it appears that universal cervical length screening and treatment with vaginal progesterone is a cost-effective strategy to prevent preterm birth and resultant neonatal morbidity and mortality. Similarly, Campbell<sup>178</sup> has concluded that, in light of the available evidence, doing nothing to prevent preterm birth is no longer an option.

### Implications for practice

The present IPD metaanalysis provides compelling evidence of the benefit of vaginal progesterone to prevent preterm birth and neonatal morbidity/mortality in women with a sonographic short cervix. Importantly, there was no evidence of demonstrable risk. This IPD metaanalysis indicates that vaginal progesterone is effective in women with and without a history of preterm birth and a short cervix. Therefore, we recommend that transvaginal sonographic measurement of cervical length be performed at 19-24 weeks of gestation. Vaginal progesterone at a dose of 90 mg/d should be considered for use in patients with a short cervix, mainly those with a cervical length between 10-20 mm, from 20-36 6/7 weeks of gestation.

### Implications for research

A properly-designed randomized controlled trial in twin gestations is needed to determine the efficacy of vaginal pro-

gesterone to prevent preterm birth and neonatal morbidity and mortality in women with a short cervix. The results of this metaanalysis suggest that this subset of twin gestations is likely to benefit from progesterone.

Another issue that deserves further investigation is the optimal management of women who have an extremely short cervix with some evidence of subclinical intraamniotic infection/inflammation (eg, patients with a short cervix and amniotic fluid sludge<sup>179</sup> or those with elevated concentrations of cervical proinflammatory cytokines<sup>180</sup> or other biomarkers for intraamniotic infection/inflammation<sup>158,159</sup>). Antibiotic and antiinflammatory agents may have a role under these circumstances. Indeed, intraamniotic infection can be eradicated in patients with a short cervix<sup>118</sup>—such observations are consistent with experimental infection in nonhuman primates.<sup>181</sup> The complexities of addressing this important clinical question should not be underestimated. ■

REFERENCES

1. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet* 2008;371:75-84.
2. Mathews TJ, Menacker F, MacDorman MF. Infant mortality statistics from the 2002 period: linked birth/infant death data set. *Natl Vital Stat Rep* 2004;53:1-29.
3. Beck S, Wojdyla D, Say L, et al. The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity. *Bull World Health Organ* 2010;88:31-8.
4. Martin JA, Hamilton BE, Sutton PD, et al. Births: final data for 2007. *Natl Vital Stat Rep* 2010;58:1-85.
5. Ananth CV, Vintzileos AM. Epidemiology of preterm birth and its clinical subtypes. *J Matern Fetal Neonatal Med* 2006;19:773-82.
6. Ananth CV, Vintzileos AM. Maternal-fetal conditions necessitating a medical intervention resulting in preterm birth. *Am J Obstet Gynecol* 2006;195:1557-63.
7. Ananth CV, Joseph KS, Demissie K, Vintzileos AM. Trends in twin preterm birth subtypes in the United States, 1989 through 2000: impact on perinatal mortality. *Am J Obstet Gynecol* 2005;193:1076-82.
8. Chauhan SP, Scardo JA, Hayes E, Abuhamad AZ, Berghella V. Twins: prevalence, problems, and preterm births. *Am J Obstet Gynecol* 2010;203:305-15.
9. Stone J, Ferrara L, Kamrath J, et al. Contemporary outcomes with the latest 1000 cases of

- multifetal pregnancy reduction (MPR). *Am J Obstet Gynecol* 2008;199:406.e1-4.
10. Luke B, Brown MB. Maternal morbidity and infant death in twin vs triplet and quadruplet pregnancies. *Am J Obstet Gynecol* 2008;198:401.e1-10.
11. Barri PN, Coroleu B, Clua E, Tur R. Prevention of prematurity by single embryo transfer. *J Perinat Med* 2011;39:237-40.
12. Pandian Z, Templeton A, Bhattacharya S. Modification of assisted reproduction techniques to prevent preterm birth. *Clin Obstet Gynecol* 2004;47:833-41; discussion 81-2.
13. Adashi EY, Ekins MN, Lacoursiere Y. On the discharge of Hippocratic obligations: challenges and opportunities. *Am J Obstet Gynecol* 2004;190:885-93.
14. Adashi EY, Barri PN, Berkowitz R, et al. Infertility therapy-associated multiple pregnancies (births): an ongoing epidemic. *Reprod Biomed Online* 2003;7:515-42.
15. Langhoff-Roos J, Kesmodel U, Jacobsson B, Rasmussen S, Vogel I. Spontaneous preterm delivery in primiparous women at low risk in Denmark: population based study. *BMJ* 2006;332:937-9.
16. Romero R. Prenatal medicine: the child is the father of the man 1996. *J Matern Fetal Neonatal Med* 2009;22:636-9.
17. Di Renzo GC. The great obstetrical syndromes. *J Matern Fetal Neonatal Med* 2009;22:633-5.
18. Romero R, Mazor M, Munoz H, Gomez R, Galasso M, Sherer DM. The preterm labor syndrome. *Ann N Y Acad Sci* 1994;734:414-29.
19. Gomez R, Romero R, Nien JK, et al. Antibiotic administration to patients with preterm premature rupture of membranes does not eradicate intra-amniotic infection. *J Matern Fetal Neonatal Med* 2007;20:167-73.
20. Ma S, Yang L, Romero R, Cui Y. Varying coefficient model for gene-environment interaction: a non-linear look. *Bioinformatics* 2011;27:2119-26.
21. Macones GA, Parry S, Elkousy M, Clothier B, Ural SH, Strauss JF III. A polymorphism in the promoter region of TNF and bacterial vaginosis: preliminary evidence of gene-environment interaction in the etiology of spontaneous preterm birth. *Am J Obstet Gynecol* 2004;190:1504-8; discussion 3A.
22. Parimi N, Tromp G, Kuivaniemi H, et al. Analytical approaches to detect maternal/fetal genotype incompatibilities that increase risk of pre-eclampsia. *BMC Med Genet* 2008;9:60.
23. Csapo AI. The 'see-saw' theory of parturition. *Ciba Found Symp* 1977;47:159-210.
24. Kerenyi T. Forgotten "father of progesterone." *Am J Obstet Gynecol* 2010;202:e10-1.
25. Csapo A. The luteo-placental shift, the guardian of pre-natal life. *Postgrad Med J* 1969;45:57-64.
26. Csapo AI, Pulkkinen MO, Wiest WG. Effects of luteectomy and progesterone replacement therapy in early pregnant patients. *Am J Obstet Gynecol* 1973;115:759-65.

27. Romero R. Prevention of spontaneous preterm birth: the role of sonographic cervical length in identifying patients who may benefit from progesterone treatment. *Ultrasound Obstet Gynecol* 2007;30:675-86.
28. Romero R, Espinoza J, Kusanovic JP, et al. The preterm parturition syndrome. *BJOG* 2006;113(Suppl):17-42.
29. Elovitz MA, Mrinalini C. The use of progestational agents for preterm birth: lessons from a mouse model. *Am J Obstet Gynecol* 2006;195:1004-10.
30. Carbone B, Dallot E, Haddad B, Ferré F, Cabrol D. Effects of progesterone on prostaglandin E(2)-induced changes in glycosaminoglycan synthesis by human cervical fibroblasts in culture. *Mol Hum Reprod* 2000;6:661-4.
31. Facchinetti F, Paganelli S, Comitini G, Dante G, Volpe A. Cervical length changes during preterm cervical ripening: effects of 17-alpha-hydroxyprogesterone caproate. *Am J Obstet Gynecol* 2007;196:453.e1-4; discussion 21.
32. Denison FC, Calder AA, Kelly RW. The action of prostaglandin E2 on the human cervix: stimulation of interleukin 8 and inhibition of secretory leukocyte protease inhibitor. *Am J Obstet Gynecol* 1999;180:614-20.
33. Kelly RW, Leask R, Calder AA. Chorionic production of interleukin-8 and mechanism of parturition. *Lancet* 1992;339:776-7.
34. Rajabi M, Solomon S, Poole AR. Hormonal regulation of interstitial collagenase in the uterine cervix of the pregnant guinea pig. *Endocrinology* 1991;128:863-71.
35. Rodriguez HA, Kass L, Varayoud J, et al. Collagen remodeling in the guinea-pig uterine cervix at term is associated with a decrease in progesterone receptor expression. *Mol Hum Reprod* 2003;9:807-13.
36. Sato T, Ito A, Mori Y, Yamashita K, Hayakawa T, Nagase H. Hormonal regulation of collagenolysis in uterine cervical fibroblasts: modulation of synthesis of procollagenase, prostromelysin and tissue inhibitor of metalloproteinases (TIMP) by progesterone and oestradiol-17 beta. *Biochem J* 1991;275:645-50.
37. Uldbjerg N, Ekman G, Malmstrom A, Olsson K, Ulfsten U. Ripening of the human uterine cervix related to changes in collagen, glycosaminoglycans, and collagenolytic activity. *Am J Obstet Gynecol* 1983;147:662-6.
38. Stjernholm Y, Sahlin L, Akerberg S, et al. Cervical ripening in humans: potential roles of estrogen, progesterone, and insulin-like growth factor-I. *Am J Obstet Gynecol* 1996;174:1065-71.
39. Naftolin F, Stubblefield P. Dilatation of the uterine cervix: connective tissue biology and clinical management. New York, NY: Raven Press Books; 1980.
40. Chwalisz K, Benson M, Scholz P, Daum J, Beier HM, Hegele-Hartung C. Cervical ripening with the cytokines interleukin 8, interleukin 1 beta and tumor necrosis factor alpha in guinea-pigs. *Hum Reprod* 1994;9:2173-81.
41. Chwalisz K, Garfield RE. Regulation of the uterus and cervix during pregnancy and labor:

- role of progesterone and nitric oxide. *Ann N Y Acad Sci* 1997;828:238-53.
- 42.** Chwalisz K, Garfield RE. Nitric oxide as the final metabolic mediator of cervical ripening. *Hum Reprod* 1998;13:245-8.
- 43.** Mahendroo MS, Cala KM, Russell DW. 5 Alpha-reduced androgens play a key role in murine parturition. *Mol Endocrinol* 1996;10:380-92.
- 44.** Mahendroo MS, Porter A, Russell DW, Word RA. The parturition defect in steroid 5 alpha-reductase type 1 knockout mice is due to impaired cervical ripening. *Mol Endocrinol* 1999;13:981-92.
- 45.** Word RA, Li XH, Hnat M, Carrick K. Dynamics of cervical remodeling during pregnancy and parturition: mechanisms and current concepts. *Semin Reprod Med* 2007;25:69-79.
- 46.** Garfield RE, Puri CP, Csapo AI. Endocrine, structural, and functional changes in the uterus during premature labor. *Am J Obstet Gynecol* 1982;142:21-7.
- 47.** Saito Y, Takahashi S, Maki M. Effects of some drugs on ripening of uterine cervix in non-pregnant castrated and pregnant rats. *Tohoku J Exp Med* 1981;133:205-20.
- 48.** Elovitz M, Wang Z. Medroxyprogesterone acetate, but not progesterone, protects against inflammation-induced parturition and intrauterine fetal demise. *Am J Obstet Gynecol* 2004;190:693-701.
- 49.** Ito A, Imada K, Sato T, Kubo T, Matsushima K, Mori Y. Suppression of interleukin 8 production by progesterone in rabbit uterine cervix. *Biochem J* 1994;301:183-6.
- 50.** Marx SG, Wentz MJ, Mackay LB, et al. Effects of progesterone on iNOS, COX-2, and collagen expression in the cervix. *J Histochem Cytochem* 2006;54:623-39.
- 51.** Stiemer B, Elger W. Cervical ripening of the rat in dependence on endocrine milieu; effects of antigestagens. *J Perinat Med* 1990;18:419-29.
- 52.** Zuidema LJ, Khan-Dawood F, Dawood MY, Work BA Jr. Hormones and cervical ripening: dehydroepiandrosterone sulfate, estradiol, estriol, and progesterone. *Am J Obstet Gynecol* 1986;155:1252-4.
- 53.** Chwalisz K, Shi Shao O, Neff G, Elger J. The effect of antigestagen ZK 98, 199 on the uterine cervix. *Acta Endocrinol* 1987;283:113.
- 54.** Elliott CL, Brennand JE, Calder AA. The effects of mifepristone on cervical ripening and labor induction in primigravidae. *Obstet Gynecol* 1998;92:804-9.
- 55.** Norman J. Antiprogestones. *Br J Hosp Med* 1991;45:372-5.
- 56.** Giacalone PL, Daures JP, Faure JM, Boulot P, Hedon B, Lafargue F. The effects of mifepristone on uterine sensitivity to oxytocin and on fetal heart rate patterns. *Eur J Obstet Gynecol Reprod Biol* 2001;97:30-4.
- 57.** Hegele-Hartung C, Chwalisz K, Beier HM, Elger W. Ripening of the uterine cervix of the guinea-pig after treatment with the progesterone antagonist onapristone (ZK 98.299): an electron microscopic study. *Hum Reprod* 1989;4:369-77.
- 58.** Stenlund PM, Ekman G, Aedo AR, Bygdeman M. Induction of labor with mifepristone—a randomized, double-blind study versus placebo. *Acta Obstet Gynecol Scand* 1999;78:793-8.
- 59.** Stys SJ, Clewell WH, Meschia G. Changes in cervical compliance at parturition independent of uterine activity. *Am J Obstet Gynecol* 1978;130:414-8.
- 60.** Wolf JP, Sinosich M, Anderson TL, Ulmann A, Baulieu EE, Hodgen GD. Progesterone antagonist (RU 486) for cervical dilation, labor induction, and delivery in monkeys: effectiveness in combination with oxytocin. *Am J Obstet Gynecol* 1989;160:45-7.
- 61.** Allan GF, Tsai SY, Tsai MJ, O'Malley BW. Ligand-dependent conformational changes in the progesterone receptor are necessary for events that follow DNA binding. *Proc Natl Acad Sci U S A* 1992;89:11750-4.
- 62.** Beato M. Gene regulation by steroid hormones. *Cell* 1989;56:335-44.
- 63.** Brosens JJ, Tullet J, Varshochi R, Lam EW. Steroid receptor action. *Best Pract Res Clin Obstet Gynaecol* 2004;18:265-83.
- 64.** Henderson D, Wilson T. Reduced binding of progesterone receptor to its nuclear response element after human labor onset. *Am J Obstet Gynecol* 2001;185:579-85.
- 65.** Merlino AA, Welsh TN, Tan H, et al. Nuclear progesterone receptors in the human pregnancy myometrium: evidence that parturition involves functional progesterone withdrawal mediated by increased expression of progesterone receptor-A. *J Clin Endocrinol Metab* 2007;92:1927-33.
- 66.** Oh SY, Kim CJ, Park I, et al. Progesterone receptor isoform (A/B) ratio of human fetal membranes increases during term parturition. *Am J Obstet Gynecol* 2005;193:1156-60.
- 67.** Power RF, Conneely OM, O'Malley BW. New insights into activation of the steroid hormone receptor superfamily. *Trends Pharmacol Sci* 1992;13:318-23.
- 68.** Cabrol D, Carbonne B, Bienkiewicz A, Dalot E, Alj AE, Cedard L. Induction of labor and cervical maturation using mifepristone (RU 486) in the late pregnant rat. Influence of a cyclooxygenase inhibitor (Diclofenac). *Prostaglandins* 1991;42:71-9.
- 69.** Tsai MJ, O'Malley BW. Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* 1994;63:451-86.
- 70.** Vegeto E, Shahbaz MM, Wen DX, Goldman ME, O'Malley BW, McDonnell DP. Human progesterone receptor A form is a cell- and promoter-specific repressor of human progesterone receptor B function. *Mol Endocrinol* 1993;7:1244-55.
- 71.** Denner LA, Weigel NL, Maxwell BL, Schrader WT, O'Malley BW. Regulation of progesterone receptor-mediated transcription by phosphorylation. *Science* 1990;250:1740-3.
- 72.** Fomin VP, Cox BE, Word RA. Effect of progesterone on intracellular Ca<sup>2+</sup> homeostasis in human myometrial smooth muscle cells. *Am J Physiol* 1999;276:C379-85.
- 73.** McEwen BS. Non-genomic and genomic effects of steroids on neural activity. *Trends Pharmacol Sci* 1991;12:141-7.
- 74.** Meizel S, Turner KO. Progesterone acts at the plasma membrane of human sperm. *Mol Cell Endocrinol* 1991;77:R1-5.
- 75.** Mesiano S, Chan EC, Fitter JT, Kwek K, Yeo G, Smith R. Progesterone withdrawal and estrogen activation in human parturition are coordinated by progesterone receptor A expression in the myometrium. *J Clin Endocrinol Metab* 2002;87:2924-30.
- 76.** Schumacher M. Rapid membrane effects of steroid hormones: an emerging concept in neuroendocrinology. *Trends Neurosci* 1990;13:359-62.
- 77.** Condon JC, Jeyasuria P, Faust JM, Wilson JW, Mendelson CR. A decline in the levels of progesterone receptor coactivators in the pregnant uterus at term may antagonize progesterone receptor function and contribute to the initiation of parturition. *Proc Natl Acad Sci USA* 2003;100:9518-23.
- 78.** Chapman NR, Kennelly MM, Harper KA, Europe-Finner GN, Robson SC. Examining the spatio-temporal expression of mRNA encoding the membrane-bound progesterone receptor-alpha isoform in human cervix and myometrium during pregnancy and labor. *Mol Hum Reprod* 2006;12:19-24.
- 79.** Madsen G, Zakar T, Ku CY, Sanborn BM, Smith R, Mesiano S. Prostaglandins differentially modulate progesterone receptor-A and -B expression in human myometrial cells: evidence for prostaglandin-induced functional progesterone withdrawal. *J Clin Endocrinol Metab* 2004;89:1010-3.
- 80.** Condon JC, Hardy DB, Kovaric K, Mendelson CR. Upregulation of the progesterone receptor (PR)-C isoform in laboring myometrium by activation of nuclear factor-kappaB may contribute to the onset of labor through inhibition of PR function. *Mol Endocrinol* 2006;20:764-75.
- 81.** Falkenstein E, Norman AW, Wehling M. Mannheim classification of nongenomically initiated (rapid) steroid action(s). *J Clin Endocrinol Metab* 2000;85:2072-5.
- 82.** Bennett P, Allport V, Loudon J, Elliott C. Prostaglandins, the fetal membranes and the cervix. *Front Horm Res* 2001;27:147-64.
- 83.** Kofinas AD, Rose JC, Koritnik DR, Meis PJ. Progesterone and estradiol concentrations in non-pregnant and pregnant human myometrium: effect of progesterone and estradiol on cyclic adenosine monophosphate-phosphodiesterase activity. *J Reprod Med* 1990;35:1045-50.
- 84.** Losel R, Wehling M. Nongenomic actions of steroid hormones. *Nat Rev Mol Cell Biol* 2003;4:46-56.
- 85.** Losel RM, Falkenstein E, Feuring M, et al. Nongenomic steroid action: controversies,

questions, and answers. *Physiol Rev* 2003;83:965-1016.

**86.** Perusquia M, Garcia-Yanez E, Ibanez R, Kubli-Garfias C. Non-genomic mechanism of action of delta-4 and 5-reduced androgens and progestins on the contractility of the isolated rat myometrium. *Life Sci* 1990;47:1547-53.

**87.** Sager G, Orbo A, Jaeger R, Engstrom C. Non-genomic effects of progestins—inhibition of cell growth and increased intracellular levels of cyclic nucleotides. *J Steroid Biochem Mol Biol* 2003;84:1-8.

**88.** Ekman-Ordeberg G, Stjernholm Y, Wang H, Stygar D, Sahlin L. Endocrine regulation of cervical ripening in humans – potential roles for gonadal steroids and insulin-like growth factor-I. *Steroids* 2003;68:837-47.

**89.** Blackmore PF, Neulen J, Lattanzio F, Beebe SJ. Cell surface-binding sites for progesterone mediate calcium uptake in human sperm. *J Biol Chem* 1991;266:18655-9.

**90.** Fernandes MS, Pierron V, Michalovich D, et al. Regulated expression of putative membrane progesterin receptor homologues in human endometrium and gestational tissues. *J Endocrinol* 2005;187:89-101.

**91.** Grazzini E, Guillon G, Mouillac B, Zingg HH. Inhibition of oxytocin receptor function by direct binding of progesterone. *Nature* 1998;392:509-12.

**92.** Haukkamaa M. High affinity progesterone binding sites of human uterine microsomal membranes. *J Steroid Biochem* 1984;20:569-73.

**93.** Karteris E, Zervou S, Pang Y, et al. Progesterone signaling in human myometrium through two novel membrane G protein-coupled receptors: potential role in functional progesterone withdrawal at term. *Mol Endocrinol* 2006;20:1519-34.

**94.** Lye SJ, Porter DG. Demonstration that progesterone 'blocks' uterine activity in the ewe in vivo by a direct action on the myometrium. *J Reprod Fertil* 1978;52:87-94.

**95.** Challis JR, Patel FA, Pomini F. Prostaglandin dehydrogenase and the initiation of labor [Review]. *J Perinat Med* 1999;27:26-34.

**96.** Modi DN, Shah C, Puri CP. Non-genomic membrane progesterone receptors on human spermatozoa. *Soc Reprod Fertil Suppl* 2007;63:515-29.

**97.** Andersson S, Minjarez D, Yost NP, Word RA. Estrogen and progesterone metabolism in the cervix during pregnancy and parturition. *J Clin Endocrinol Metab* 2008;93:2366-74.

**98.** Yellon SM, Ebner CA, Elovitz MA. Medroxyprogesterone acetate modulates remodeling, immune cell census, and nerve fibers in the cervix of a mouse model for inflammation-induced preterm birth. *Reprod Sci* 2009;16:257-64.

**99.** Cicinelli E, de Ziegler D. Transvaginal progesterone: evidence for a new functional 'portal system' flowing from the vagina to the uterus. *Hum Reprod Update* 1999;5:365-72.

**100.** Romero R, Scoccia B, Mazor M, Wu YK, Benveniste R. Evidence for a local change in the progesterone/estrogen ratio in human parturi-

tion at term. *Am J Obstet Gynecol* 1988;159:657-60.

**101.** Mitchell BF, Mitchell JM, Chowdhury J, Tougas M, Engelen SM, Senff N, et al. Metabolites of progesterone and the pregnane X receptor: a novel pathway regulating uterine contractility in pregnancy? *Am J Obstet Gynecol* 2005;192:1304-13; discussion 1313-5. Erratum in: *Am J Obstet Gynecol* 2005;193(3 Pt 1):899.

**102.** Andersen HF, Nugent CE, Wanty SD, Hayashi RH. Prediction of risk for preterm delivery by ultrasonographic measurement of cervical length. *Am J Obstet Gynecol* 1990;163:859-67.

**103.** Iams JD, Goldenberg RL, Meis PJ, et al. The length of the cervix and the risk of spontaneous premature delivery. *N Engl J Med* 1996;334:567-72.

**104.** Hassan SS, Romero R, Berry SM, et al. Patients with an ultrasonographic cervical length < or = 15 mm have nearly a 50% risk of early spontaneous preterm delivery. *Am J Obstet Gynecol* 2000;182:1458-67.

**105.** To MS, Skentou C, Liao AW, Cacho A, Nicolaides KH. Cervical length and funneling at 23 weeks of gestation in the prediction of spontaneous early preterm delivery. *Ultrasound Obstet Gynecol* 2001;18:200-3.

**106.** Celik E, To M, Gajewska K, Smith GC, Nicolaides KH. Cervical length and obstetric history predict spontaneous preterm birth: development and validation of a model to provide individualized risk assessment. *Ultrasound Obstet Gynecol* 2008;31:549-54.

**107.** Crane JM, Hutchens D. Transvaginal sonographic measurement of cervical length to predict preterm birth in asymptomatic women at increased risk: a systematic review. *Ultrasound Obstet Gynecol* 2008;31:579-87.

**108.** Conde-Agudelo A, Romero R, Hassan SS, Yeo L. Transvaginal sonographic cervical length for the prediction of spontaneous preterm birth in twin pregnancies: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2010;203:128.e1-12.

**109.** Vaisbuch E, Romero R, Erez O, et al. Clinical significance of early (<20 weeks) vs late (20-24 weeks) detection of sonographic short cervix in asymptomatic women in the mid-trimester. *Ultrasound Obstet Gynecol* 2010;36:471-81.

**110.** Heath VC, Southall TR, Souka AP, Elisseou A, Nicolaides KH. Cervical length at 23 weeks of gestation: prediction of spontaneous preterm delivery. *Ultrasound Obstet Gynecol* 1998;12:312-7.

**111.** Taipale P, Hiilesmaa V. Sonographic measurement of uterine cervix at 18-22 weeks' gestation and the risk of preterm delivery. *Obstet Gynecol* 1998;92:902-7.

**112.** Heath VC, Daskalakis G, Zagaliki A, Carvalho M, Nicolaides KH. Cervicovaginal fibronectin and cervical length at 23 weeks of gestation: relative risk of early preterm delivery. *BJOG* 2000;107:1276-81.

**113.** To MS, Alfirevic Z, Heath VC, et al. Cervical cerclage for prevention of preterm delivery in women with short cervix: randomized controlled trial. *Lancet* 2004;363:1849-53.

**114.** To MS, Palaniappan V, Skentou C, Gibb D, Nicolaides KH. Elective cerclage vs ultrasound-indicated cerclage in high-risk pregnancies. *Ultrasound Obstet Gynecol* 2002;19:475-7.

**115.** Alfirevic Z, Allen-Coward H, Molina F, Vinuesa CP, Nicolaides K. Targeted therapy for threatened preterm labor based on sonographic measurement of the cervical length: a randomized controlled trial. *Ultrasound Obstet Gynecol* 2007;29:47-50.

**116.** Gomez R, Romero R, Medina L, et al. Cervicovaginal fibronectin improves the prediction of preterm delivery based on sonographic cervical length in patients with preterm uterine contractions and intact membranes. *Am J Obstet Gynecol* 2005;192:350-9.

**117.** Gomez R, Romero R, Nien JK, et al. A short cervix in women with preterm labor and intact membranes: a risk factor for microbial invasion of the amniotic cavity. *Am J Obstet Gynecol* 2005;192:678-89.

**118.** Hassan S, Romero R, Hendler I, et al. A sonographic short cervix as the only clinical manifestation of intra-amniotic infection. *J Perinat Med* 2006;34:13-9.

**119.** Kusanovic JP, Espinoza J, Romero R, et al. Clinical significance of the presence of amniotic fluid 'sludge' in asymptomatic patients at high risk for spontaneous preterm delivery. *Ultrasound Obstet Gynecol* 2007;30:706-14.

**120.** Palma-Dias RS, Fonseca MM, Stein NR, Schmidt AP, Magalhaes JA. Relation of cervical length at 22-24 weeks of gestation to demographic characteristics and obstetric history. *Braz J Med Biol Res* 2004;37:737-44.

**121.** Tekesin I, Eberhart LH, Schaefer V, Wallwiener D, Schmidt S. Evaluation and validation of a new risk score (CLEOPATRA score) to predict the probability of premature delivery for patients with threatened preterm labor. *Ultrasound Obstet Gynecol* 2005;26:699-706.

**122.** To MS, Skentou CA, Royston P, Yu CK, Nicolaides KH. Prediction of patient-specific risk of early preterm delivery using maternal history and sonographic measurement of cervical length: a population-based prospective study. *Ultrasound Obstet Gynecol* 2006;27:362-7.

**123.** Owen J, Yost N, Berghella V, et al. Mid-trimester endovaginal sonography in women at high risk for spontaneous preterm birth. *JAMA* 2001;286:1340-8.

**124.** Keirse MJ. Progesterone and preterm: seventy years of "deja vu" or "still to be seen"? *Birth* 2004;31:230-5.

**125.** da Fonseca EB, Bittar RE, Carvalho MH, Zugaib M. Prophylactic administration of progesterone by vaginal suppository to reduce the incidence of spontaneous preterm birth in women at increased risk: a randomized placebo-controlled double-blind study. *Am J Obstet Gynecol* 2003;188:419-24.

- 126.** Hartikainen-Sorri AL, Kaupilla A, Tuimala R. Inefficacy of 17 alpha-hydroxyprogesterone caproate in the prevention of prematurity in twin pregnancy. *Obstet Gynecol* 1980;56:692-5.
- 127.** Hauth JC, Gilstrap LC III, Brekken AL, Hauth JM. The effect of 17 alpha-hydroxyprogesterone caproate on pregnancy outcome in an active-duty military population. *Am J Obstet Gynecol* 1983;146:187-90.
- 128.** Johnson JW, Austin KL, Jones GS, Davis GH, King TM. Efficacy of 17alpha-hydroxyprogesterone caproate in the prevention of premature labor. *N Engl J Med* 1975;293:675-80.
- 129.** Meis PJ, Klebanoff M, Thom E, et al. Prevention of recurrent preterm delivery by 17 alpha-hydroxyprogesterone caproate. *N Engl J Med* 2003;348:2379-85.
- 130.** Yemini M, Borenstein R, Drazan E, et al. Prevention of premature labor by 17 alpha-hydroxyprogesterone caproate. *Am J Obstet Gynecol* 1985;151:574-7.
- 131.** Fonseca EB, Celik E, Parra M, Singh M, Nicolaides KH. Progesterone and the risk of preterm birth among women with a short cervix. *N Engl J Med* 2007;357:462-9.
- 132.** DeFranco EA, O'Brien JM, Adair CD, et al. Vaginal progesterone is associated with a decrease in risk for early preterm birth and improved neonatal outcome in women with a short cervix: a secondary analysis from a randomized, double-blind, placebo-controlled trial. *Ultrasound Obstet Gynecol* 2007;30:697-705.
- 133.** Hassan SS, Romero R, Vidyadhari D, et al. Vaginal progesterone reduces the rate of preterm birth in women with a sonographic short cervix: a multicenter, randomized, double-blind, placebo-controlled trial. *Ultrasound Obstet Gynecol* 2011;38:18-31.
- 134.** Rode L, Klein K, Nicolaides K, Krampl-Bettelheim E, Tabor A. Prevention of preterm delivery in twin gestations (PREDICT): a multicenter randomized placebo-controlled trial on the effect of vaginal micronized progesterone. *Ultrasound Obstet Gynecol* 2011;38:272-80.
- 135.** Klein K, Rode L, Nicolaides K, Krampl-Bettelheim E, Tabor A. Vaginal micronized progesterone and the risk of preterm delivery in high-risk twin pregnancies—secondary analysis of a placebo-controlled randomized trial and meta-analysis. *Ultrasound Obstet Gynecol* 2011;38:281-7.
- 136.** Cetingoz E, Cam C, Sakalli M, Karateke A, Celik C, Sancak A. Progesterone effects on preterm birth in high-risk pregnancies: a randomized placebo-controlled trial. *Arch Gynecol Obstet* 2011;283:423-9.
- 137.** Higgins JPT, Altman DG, Sterne JAC. Chapter 8: assessing risk of bias in included studies. In: Higgins JPT, Green S, eds. *Cochrane handbook for systematic reviews of interventions*, version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011.
- 138.** Stewart LA, Parmar MK. Meta-analysis of the literature or of individual patient data: is there a difference? *Lancet* 1993;341:418-22.
- 139.** Thompson SG, Higgins JP. Treating individuals 4: can meta-analysis help target interventions at individuals most likely to benefit? *Lancet* 2005;365:341-6.
- 140.** Sutton AJ, Kendrick D, Coupland CA. Meta-analysis of individual- and aggregate-level data. *Stat Med* 2008;27:651-69.
- 141.** Riley RD, Lambert PC, Abo-Zaid G. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ* 2010;340:c221.
- 142.** Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
- 143.** Honest H, Bachmann LM, Coomarasamy A, Gupta JK, Kleijnen J, Khan KS. Accuracy of cervical transvaginal sonography in predicting preterm birth: a systematic review. *Ultrasound Obstet Gynecol* 2003;22:305-22.
- 144.** O'Brien JM, Adair CD, Lewis DF, et al. Progesterone vaginal gel for the reduction of recurrent preterm birth: primary results from a randomized, double-blind, placebo-controlled trial. *Ultrasound Obstet Gynecol* 2007;30:687-96.
- 145.** Simmonds MC, Higgins JP, Stewart LA, Tierney JF, Clarke MJ, Thompson SG. Meta-analysis of individual patient data from randomized trials: a review of methods used in practice. *Clin Trials* 2005;2:209-17.
- 146.** Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
- 147.** Altman DG. Confidence intervals for the number needed to treat. *BMJ* 1998;317:1309-12.
- 148.** Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
- 149.** Rothwell PM. Treating individuals 2. Subgroup analysis in randomized controlled trials: importance, indications, and interpretation. *Lancet* 2005;365:176-86.
- 150.** Klebanoff MA. Subgroup analysis in obstetrics clinical trials. *Am J Obstet Gynecol* 2007;197:119-22.
- 151.** Berlin JA, Santanna J, Schmid CH, Szczech LA, Feldman HI. Individual patient- versus group-level data meta-regressions for the investigation of treatment effect modifiers: ecological bias rears its ugly head. *Stat Med* 2002;21:371-87.
- 152.** Gates S, Brocklehurst P. How should randomized trials including multiple pregnancies be analyzed? *BJOG* 2004;111:213-9.
- 153.** Borna S, Sahabi N. Progesterone for maintenance tocolytic therapy after threatened preterm labor: a randomized controlled trial. *Aust N Z J Obstet Gynaecol* 2008;48:58-63.
- 154.** Majhi P, Bagga R, Kalra J, Sharma M. Intravaginal use of natural micronized progesterone to prevent pre-term birth: a randomized trial in India. *J Obstet Gynaecol* 2009;29:493-8.
- 155.** Norman JE, Mackenzie F, Owen P, et al. Progesterone for the prevention of preterm birth in twin pregnancy (STOPPIT): a randomized, double-blind, placebo-controlled study and meta-analysis. *Lancet* 2009;373:2034-40.
- 156.** Sharami SH, Zahiri Z, Shakiba M, Milani F. Maintenance therapy by vaginal progesterone after threatened idiopathic preterm labor: a randomized placebo-controlled double-blind trial. *Int J Fertil Steril* 2010;4:45-50.
- 157.** O'Brien JM, Steichen JJ, Phillips JA, Creasy GW. Two year infant outcomes for children exposed to supplemental intravaginal progesterone gel in utero: secondary analysis of a multicenter, randomized, double-blind, placebo-controlled trial. *Am J Obstet Gynecol* 2012;206(S1):S223.
- 158.** Kiefer DG, Keeler SM, Rust OA, Wayock CP, Vintzileos AM, Hanna N. Is midtrimester short cervix a sign of intraamniotic inflammation? *Am J Obstet Gynecol* 2009;200:374.e1-5.
- 159.** Vaisbuch E, Hassan SS, Mazaki-Tovi S, et al. Patients with an asymptomatic short cervix (< or =15 mm) have a high rate of subclinical intraamniotic inflammation: implications for patient counseling. *Am J Obstet Gynecol* 2010;202:433.e1-8.
- 160.** Lockwood C. The real progesterone story. *Contemp OB/GYN* 2011;56:10-5.
- 161.** Berghella V, Rafael TJ, Szychowski JM, Rust OA, Owen J. Cerclage for short cervix on ultrasonography in women with singleton gestations and previous preterm birth: a meta-analysis. *Obstet Gynecol* 2011;117:663-71.
- 162.** Gupta M, Emary K, Impey L. Emergency cervical cerclage: predictors of success. *J Matern Fetal Neonatal Med* 2010;23:670-4.
- 163.** Owen J, Hankins G, Iams JD, et al. Multicenter randomized trial of cerclage for preterm birth prevention in high-risk women with shortened midtrimester cervical length. *Am J Obstet Gynecol* 2009;201:375.e1-8.
- 164.** Fox NS, Chervenak FA. Cervical cerclage: a review of the evidence. *Obstet Gynecol Surv* 2008;63:58-65.
- 165.** Vidaeff AC, Ramin SM. Management strategies for the prevention of preterm birth: part II—update on cervical cerclage. *Curr Opin Obstet Gynecol* 2009;21:485-90.
- 166.** Abenheim HA, Tulandi T. Cervical insufficiency: re-evaluating the prophylactic cervical cerclage. *J Matern Fetal Neonatal Med* 2009;22:510-6.
- 167.** Zaveri V, Aghajafari F, Amankwah K, Hannah M. Abdominal versus vaginal cerclage after a failed transvaginal cerclage: a systematic review. *Am J Obstet Gynecol* 2002;187:868-72.
- 168.** Romero R, Espinoza J, Erez O, Hassan S. The role of cervical cerclage in obstetric practice: can the patient who could benefit from this procedure be identified? *Am J Obstet Gynecol* 2006;194:1-9.
- 169.** Hassan SS, Romero R, Maymon E, et al. Does cervical cerclage prevent preterm delivery in patients with a short cervix? *Am J Obstet Gynecol* 2001;184:1325-9.
- 170.** Althuisius SM, Dekker GA, Hummel P, van Geijn HP. Cervical incompetence prevention randomized cerclage trial: emergency cerclage

with bed rest versus bed rest alone. *Am J Obstet Gynecol* 2003;189:907-10.

**171.** Althuisius SM, Dekker GA, van Geijn HP, Bekedam DJ, Hummel P. Cervical incompetence prevention randomized cerclage trial (CIPRACT): study design and preliminary results. *Am J Obstet Gynecol* 2000;183:823-9.

**172.** Rouse DJ, Caritis SN, Peaceman AM, et al. A trial of 17 alpha-hydroxyprogesterone caproate to prevent prematurity in twins. *N Engl J Med* 2007;357:454-61.

**173.** Briery CM, Veillon EW, Klauser CK, et al. Progesterone does not prevent preterm births in women with twins. *South Med J* 2009;102:900-4.

**174.** Combs CA, Garite T, Maurel K, Das A, Porto M. 17-Hydroxyprogesterone caproate for twin pregnancy: a double-blind, randomized

clinical trial. *Am J Obstet Gynecol* 2011;204:211.e1-8.

**175.** Romero R, Gomez R, Mazor M, Ghezzi F, Yoon BH. The preterm labor syndrome. In: Elder M, Romero R, Lamont R, eds. *Preterm labor*. New York, NY: Churchill Livingstone; 1997:29-49.

**176.** Cahill AG, Odibo AO, Caughey AB, et al. Universal cervical length screening and treatment with vaginal progesterone to prevent preterm birth: a decision and economic analysis. *Am J Obstet Gynecol* 2010;202:548.e1-8.

**177.** Werner EF, Han CS, Pettker CM, et al. Universal cervical length screening to prevent preterm birth: a cost-effectiveness analysis. *Ultrasound Obstet Gynecol* 2011;38:32-7.

**178.** Campbell S. Universal cervical-length screening and vaginal progesterone prevents

early preterm births, reduces neonatal morbidity and is cost saving: doing nothing is no longer an option. *Ultrasound Obstet Gynecol* 2011;38:1-9.

**179.** Romero R, Schaudinn C, Kusanovic JP, et al. Detection of a microbial biofilm in intraamniotic infection. *Am J Obstet Gynecol* 2008;198:135.e1-5.

**180.** Sakai M, Shiozaki A, Tabata M, et al. Evaluation of effectiveness of prophylactic cerclage of a short cervix according to interleukin-8 in cervical mucus. *Am J Obstet Gynecol* 2006;194:14-9.

**181.** Gravett MG, Adams KM, Sadowsky DW, et al. Immunomodulators plus antibiotics delay preterm delivery after experimental intraamniotic infection in a nonhuman primate model. *Am J Obstet Gynecol* 2007;197:518.e1-8.